

ANALYSIS OF THE $\frac{3}{2}^{\pm}$ PENTAQUARK STATES IN THE DIQUARK-DIQUARK-ANTIQUARK MODEL WITH QCD SUM RULES

Zhi-Gang Wang ¹

Department of Physics, North China Electric Power University, Baoding 071003, P. R. China

Abstract

In this article, we construct the axialvector-diquark-scalar-diquark-antiquark type and the axialvector-diquark-axialvector-diquark-antiquark type interpolating currents, and study the $J^P = \frac{3}{2}^{\pm}$ hidden-charm pentaquark states with the strangeness $S = 0, -1, -2, -3$ systematically using the QCD sum rules. The predicted masses of the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$ are compatible with the experimental value $M_{P_c(4380)} = 4380 \pm 8 \pm 29$ MeV from the LHCb collaboration, more experimental data are still needed to identify the $P_c(4380)$ unambiguously.

PACS number: 12.39.Mk, 14.20.Lq, 12.38.Lg

Key words: Pentaquark states, QCD sum rules

1 Introduction

In 2015, the LHCb collaboration studied the $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays with the data sample corresponds to an integrated luminosity of 3 fb^{-1} acquired with the LHCb detector from 7 and 8 TeV pp collisions, and observed two pentaquark candidates $P_c(4380)$ and $P_c(4450)$ in the $J/\psi p$ invariant mass distributions with the significances of more than 9 standard deviations [1]. They performed the amplitude analysis on all relevant masses and decay angles of the six-dimensional data using the helicity formalism and Breit-Wigner amplitudes to describe all resonances. The Breit-Wigner masses and widths are $M_{P_c(4380)} = 4380 \pm 8 \pm 29$ MeV, $M_{P_c(4450)} = 4449.8 \pm 1.7 \pm 2.5$ MeV, $\Gamma_{P_c(4380)} = 205 \pm 18 \pm 86$ MeV, and $\Gamma_{P_c(4450)} = 39 \pm 5 \pm 19$ MeV, respectively. The preferred quantum numbers of the $P_c(4380)$ and $P_c(4450)$ are $J^P = \frac{3}{2}^{-}$ and $\frac{5}{2}^{+}$, respectively.

Recently, the LHCb collaboration inspected the $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays for the presence of $J/\psi p$ or $J/\psi K^-$ contributions with minimal assumptions about $K^- p$ contributions (such as their number, their resonant or nonresonant nature, or their lineshapes), and observed that at more than 9 standard deviations the $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays cannot be described with the $K^- p$ contributions alone, and the $J/\psi p$ contributions play a dominant role in this incompatibility, and obtained model-independent support for the evidences of the $P_c^+(4380/4500) \rightarrow J/\psi p$ decays [2].

Furthermore, the LHCb collaboration performed a full amplitude analysis of the $\Lambda_b^0 \rightarrow J/\psi \pi^- p$ decays allowing for previously observed conventional ($p\pi^-$) and exotic ($J/\psi p$ and $J/\psi \pi^-$) resonances, and observed that a significantly better description of the data can be achieved by either including the two $P_c^+(4380/4450)$ states or the $Z_c^-(4200)$ state [3].

There have been several possible assignments since the observations of the $P_c(4380)$ and $P_c(4450)$, such as the molecule-like pentaquark states [4] (or not the molecular pentaquark states [5]), the diquark-triquark type pentaquark states [6], the diquark-diquark-antiquark type pentaquark states [7, 8, 9, 10, 11], re-scattering effects [12], etc. We can diagnose their resonant or non-resonant nature using photoproduction (or pionproduction) off a proton target [13]. The photoproduction process does not satisfy the so-called anomalous triangle singularity condition, their presence in the J/ψ photoproduction is crucial to conforming the resonant nature of such states as opposed to their being kinematical effects.

The QCD sum rules is a powerful theoretical tool in studying the ground state heavy baryon states [14, 15, 16, 17]. We usually take the diquarks as the basic constituents in constructing the baryon currents and pentaquark currents. The attractive interactions induced by one-gluon

¹E-mail: zgwang@aliyun.com.

exchange favor formation of the diquark states in color antitriplet, flavor antitriplet and spin singlet $\varepsilon^{ijk} q_j^T C \gamma_5 q_k'$ or in color antitriplet, flavor sextet and spin triplet $\varepsilon^{ijk} q_j^T C \gamma_\mu q_k'$, where the i, j and k are color indexes [18], the QCD sum rules also favor formation of the diquark states $\varepsilon^{ijk} q_j^T C \gamma_5 q_k'$ and $\varepsilon^{ijk} q_j^T C \gamma_\mu q_k'$ [19, 20], so we can take the scalar or axialvector diquark states as the basic constituents in constructing currents to interpolate the lowest pentaquark states. In our previous works, we observed that the heavy-light scalar and axialvector diquark states have almost degenerate masses [19], the masses of the light axialvector diquark states lie (150–200) MeV above that of the corresponding light scalar diquark states [20] from the QCD sum rules. We expect that the ground state diquark-diquark-antiquark type hidden-charm pentaquark states consist of a light scalar (or axialvector) diquark, a charm scalar (or axialvector) diquark and an anti-charm-quark.

In Refs.[9, 10], we take the light scalar diquark states, heavy axialvector diquark states and heavy scalar diquark states as the basic constituents, construct the scalar-diquark-axialvector-diquark-antiquark type and scalar-diquark-scalar-diquark-antiquark type currents to study the $J^P = \frac{3}{2}^-, \frac{5}{2}^+$ and $\frac{1}{2}^\pm$ hidden-charm pentaquark states using the QCD sum rules. The predicted masses $M_{P_c(\frac{3}{2}^-)} = 4.38 \pm 0.13 \text{ GeV}$ and $M_{P_c(\frac{5}{2}^+)} = 4.44 \pm 0.14 \text{ GeV}$ support assigning the $P_c(4380)$ and $P_c(4450)$ to be the $\frac{3}{2}^-$ and $\frac{5}{2}^+$ hidden-charm pentaquark states, respectively [9]. In Ref.[9], we observe that the empirical energy scale formula, $\mu = \sqrt{M_{X/Y/Z}^2 - (2\mathbb{M}_Q)^2}$ with the effective heavy quark mass $\mathbb{M}_Q$, in determining the ideal energy scales of the QCD spectral densities in the QCD sum rules for the tetraquark states X, Y and Z [21], can be successfully applied to study the hidden-charm pentaquark states with a slight modification $\mu = \sqrt{M_P^2 - (2\mathbb{M}_c)^2}$. In Ref.[11], we take the light axialvector diquark states, heavy axialvector diquark states and heavy scalar diquark states as the basic constituents, construct the axialvector-diquark-axialvector-diquark-antiquark type and axialvector-diquark-scalar-diquark-antiquark type interpolating currents to study the $J^P = \frac{1}{2}^\pm$ hidden-charm pentaquark states with the QCD sum rules in a systematic way.

In this article, we resort to the same routine as that in Refs.[9, 10, 11] to study the masses and pole residues of the $J^P = \frac{3}{2}^\pm$ hidden-charm pentaquark states with the QCD sum rules by constructing the axialvector-diquark-scalar-diquark-antiquark type and axialvector-diquark-axialvector-diquark-antiquark type interpolating currents, and revisit the assignment of the $P_c(4380)$.

The article is arranged as follows: we derive the QCD sum rules for the masses and pole residues of the $\frac{3}{2}^\pm$ hidden-charm pentaquark states in Sect.2; in Sect.3, we present the numerical results and discussions; and Sect.4 is reserved for our conclusion.

2 QCD sum rules for the $\frac{3}{2}^\pm$ hidden charm pentaquark states

Firstly, let us write down the two-point correlation functions $\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p)$ in the QCD sum rules,

$$\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p) = i \int d^4x e^{ip \cdot x} \langle 0 | T \{ J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(x) \bar{J}_{q_1 q_2 q_3, \nu}^{j_L j_H j}(0) \} | 0 \rangle, \quad (1)$$

where the $J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(x)$ are currents to interpolate the $J^P = \frac{3}{2}^\pm$ hidden-charm tetraquark states, the superscripts j_L and j_H are the spins of the light diquark state and the charm (or heavy) diquark state, respectively, $\vec{j} = \vec{j}_H + \vec{j}_{\bar{c}}$, the $j_{\bar{c}}$ is the spin of the charm (or heavy) antiquark, the subscripts q_1, q_2, q_3 are the light quark constituents u, d or s . Now we write down the interpolating currents

$J_{q_1 q_2 q_3, \mu}^{jLjHj}(x)$ explicitly,

$$\begin{aligned}
J_{uu\bar{u}, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} u_j^T(x) C \gamma_\mu u_k(x) u_m^T(x) C \gamma_5 c_n(x) C \bar{c}_a^T(x), \\
J_{uud, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu u_k(x) d_m^T(x) C \gamma_5 c_n(x) + 2u_j^T(x) C \gamma_\mu d_k(x) u_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \\
J_{udd, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x) C \gamma_\mu d_k(x) u_m^T(x) C \gamma_5 c_n(x) + 2d_j^T(x) C \gamma_\mu u_k(x) d_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \\
J_{ddd, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} d_j^T(x) C \gamma_\mu d_k(x) d_m^T(x) C \gamma_5 c_n(x) C \bar{c}_a^T(x), \tag{2}
\end{aligned}$$

$$\begin{aligned}
J_{uus, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu u_k(x) s_m^T(x) C \gamma_5 c_n(x) + 2u_j^T(x) C \gamma_\mu s_k(x) u_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \\
J_{uds, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu d_k(x) s_m^T(x) C \gamma_5 c_n(x) + u_j^T(x) C \gamma_\mu s_k(x) d_m^T(x) C \gamma_5 c_n(x) \\
&\quad + d_j^T(x) C \gamma_\mu s_k(x) u_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \\
J_{dds, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x) C \gamma_\mu d_k(x) s_m^T(x) C \gamma_5 c_n(x) + 2d_j^T(x) C \gamma_\mu s_k(x) d_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \tag{3}
\end{aligned}$$

$$\begin{aligned}
J_{uss, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x) C \gamma_\mu s_k(x) u_m^T(x) C \gamma_5 c_n(x) + 2s_j^T(x) C \gamma_\mu u_k(x) s_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \\
J_{dss, \mu}^{10\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x) C \gamma_\mu s_k(x) d_m^T(x) C \gamma_5 c_n(x) + 2s_j^T(x) C \gamma_\mu d_k(x) s_m^T(x) C \gamma_5 c_n(x)] C \bar{c}_a^T(x), \tag{4}
\end{aligned}$$

$$J_{sss, \mu}^{10\frac{1}{2}}(x) = \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} s_j^T(x) C \gamma_\mu s_k(x) s_m^T(x) C \gamma_5 c_n(x) C \bar{c}_a^T(x), \tag{5}$$

$$\begin{aligned}
J_{uu\bar{u}, \mu}^{11\frac{1}{2}}(x) &= \varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn} u_j^T(x) C \gamma_\mu u_k(x) u_m^T(x) C \gamma_\alpha c_n(x) \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \\
J_{uud, \mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu u_k(x) d_m^T(x) C \gamma_\alpha c_n(x) + 2u_j^T(x) C \gamma_\mu d_k(x) u_m^T(x) C \gamma_\alpha c_n(x)] \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \\
J_{udd, \mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x) C \gamma_\mu d_k(x) u_m^T(x) C \gamma_\alpha c_n(x) + 2d_j^T(x) C \gamma_\mu u_k(x) d_m^T(x) C \gamma_\alpha c_n(x)] \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \\
J_{ddd, \mu}^{11\frac{1}{2}}(x) &= \varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn} d_j^T(x) C \gamma_\mu d_k(x) d_m^T(x) C \gamma_\alpha c_n(x) \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \tag{6}
\end{aligned}$$

$$\begin{aligned}
J_{uus, \mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu u_k(x) s_m^T(x) C \gamma_\alpha c_n(x) + 2u_j^T(x) C \gamma_\mu s_k(x) u_m^T(x) C \gamma_\alpha c_n(x)] \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \\
J_{uds, \mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x) C \gamma_\mu d_k(x) s_m^T(x) C \gamma_\alpha c_n(x) + u_j^T(x) C \gamma_\mu s_k(x) d_m^T(x) C \gamma_\alpha c_n(x) \\
&\quad + d_j^T(x) C \gamma_\mu s_k(x) u_m^T(x) C \gamma_\alpha c_n(x)] \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \\
J_{dds, \mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x) C \gamma_\mu d_k(x) s_m^T(x) C \gamma_\alpha c_n(x) + 2d_j^T(x) C \gamma_\mu s_k(x) d_m^T(x) C \gamma_\alpha c_n(x)] \gamma_5 \gamma^\alpha C \bar{c}_a^T(x), \tag{7}
\end{aligned}$$

$$\begin{aligned}
J_{uss,\mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x)C\gamma_\mu s_k(x)u_m^T(x)C\gamma_\alpha c_n(x) + 2s_j^T(x)C\gamma_\mu u_k(x)s_m^T(x)C\gamma_\alpha c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{dss,\mu}^{11\frac{1}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x)C\gamma_\mu s_k(x)d_m^T(x)C\gamma_\alpha c_n(x) + 2s_j^T(x)C\gamma_\mu d_k(x)s_m^T(x)C\gamma_\alpha c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x),
\end{aligned} \tag{8}$$

$$J_{sss,\mu}^{11\frac{1}{2}}(x) = \varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}s_j^T(x)C\gamma_\mu s_k(x)s_m^T(x)C\gamma_\alpha c_n(x)\gamma_5\gamma^\alpha C\bar{c}_a^T(x), \tag{9}$$

$$\begin{aligned}
J_{uuu,\mu}^{11\frac{3}{2}}(x) &= \varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}u_j^T(x)C\gamma_\alpha u_k(x)u_m^T(x)C\gamma_\mu c_n(x)\gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{uud,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x)C\gamma_\alpha u_k(x)d_m^T(x)C\gamma_\mu c_n(x) + 2u_j^T(x)C\gamma_\alpha d_k(x)u_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{udd,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x)C\gamma_\alpha d_k(x)u_m^T(x)C\gamma_\mu c_n(x) + 2d_j^T(x)C\gamma_\alpha u_k(x)d_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{ddd,\mu}^{11\frac{3}{2}}(x) &= \varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}d_j^T(x)C\gamma_\alpha d_k(x)d_m^T(x)C\gamma_\mu c_n(x)\gamma_5\gamma^\alpha C\bar{c}_a^T(x),
\end{aligned} \tag{10}$$

$$\begin{aligned}
J_{uus,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x)C\gamma_\alpha u_k(x)s_m^T(x)C\gamma_\mu c_n(x) + 2u_j^T(x)C\gamma_\alpha s_k(x)u_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{uds,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [u_j^T(x)C\gamma_\alpha d_k(x)s_m^T(x)C\gamma_\mu c_n(x) + u_j^T(x)C\gamma_\alpha s_k(x)d_m^T(x)C\gamma_\mu c_n(x) \\
&\quad + d_j^T(x)C\gamma_\alpha s_k(x)u_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{dds,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [d_j^T(x)C\gamma_\alpha d_k(x)s_m^T(x)C\gamma_\mu c_n(x) + 2d_j^T(x)C\gamma_\alpha s_k(x)d_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x),
\end{aligned} \tag{11}$$

$$\begin{aligned}
J_{uss,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x)C\gamma_\alpha s_k(x)u_m^T(x)C\gamma_\mu c_n(x) + 2s_j^T(x)C\gamma_\alpha u_k(x)s_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x), \\
J_{dss,\mu}^{11\frac{3}{2}}(x) &= \frac{\varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{3}} [s_j^T(x)C\gamma_\alpha s_k(x)d_m^T(x)C\gamma_\mu c_n(x) + 2s_j^T(x)C\gamma_\alpha d_k(x)s_m^T(x)C\gamma_\mu c_n(x)] \gamma_5\gamma^\alpha C\bar{c}_a^T(x),
\end{aligned} \tag{12}$$

$$J_{sss,\mu}^{11\frac{3}{2}}(x) = \varepsilon^{ila}\varepsilon^{ijk}\varepsilon^{lmn}s_j^T(x)C\gamma_\alpha s_k(x)s_m^T(x)C\gamma_\mu c_n(x)\gamma_5\gamma^\alpha C\bar{c}_a^T(x), \tag{13}$$

where the i, j, k, l, m, n and a are color indices, the C is the charge conjugation matrix.

We take the isospin limit by assuming the u and d quarks have the degenerate masses, and classify the above thirty currents couple to the hidden-charm pentaquark states with degenerate

masses into the following twelve types,

1. $J_{uuu,\mu}^{10\frac{1}{2}}(x), J_{uud,\mu}^{10\frac{1}{2}}(x), J_{udd,\mu}^{10\frac{1}{2}}(x), J_{ddd,\mu}^{10\frac{1}{2}}(x);$
2. $J_{uus,\mu}^{10\frac{1}{2}}(x), J_{uds,\mu}^{10\frac{1}{2}}(x), J_{dds,\mu}^{10\frac{1}{2}}(x);$
3. $J_{uss,\mu}^{10\frac{1}{2}}(x), J_{dss,\mu}^{10\frac{1}{2}}(x);$
4. $J_{sss,\mu}^{10\frac{1}{2}}(x);$
5. $J_{uuu,\mu}^{11\frac{1}{2}}(x), J_{uud,\mu}^{11\frac{1}{2}}(x), J_{udd,\mu}^{11\frac{1}{2}}(x), J_{ddd,\mu}^{11\frac{1}{2}}(x);$
6. $J_{uus,\mu}^{11\frac{1}{2}}(x), J_{uds,\mu}^{11\frac{1}{2}}(x), J_{dds,\mu}^{11\frac{1}{2}}(x);$
7. $J_{uss,\mu}^{11\frac{1}{2}}(x), J_{dss,\mu}^{11\frac{1}{2}}(x);$
8. $J_{sss,\mu}^{11\frac{1}{2}}(x);$
9. $J_{uuu,\mu}^{11\frac{3}{2}}(x), J_{uud,\mu}^{11\frac{3}{2}}(x), J_{udd,\mu}^{11\frac{3}{2}}(x), J_{ddd,\mu}^{11\frac{3}{2}}(x);$
10. $J_{uus,\mu}^{11\frac{3}{2}}(x), J_{uds,\mu}^{11\frac{3}{2}}(x), J_{dds,\mu}^{11\frac{3}{2}}(x);$
11. $J_{uss,\mu}^{11\frac{3}{2}}(x), J_{dss,\mu}^{11\frac{3}{2}}(x);$
12. $J_{sss,\mu}^{11\frac{3}{2}}(x).$

(14)

In calculations, we choose the first current in each type for simplicity.

The currents $J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(0)$ couple potentially to the $J^P = \frac{3}{2}^\pm$ and $J^P = \frac{1}{2}^\pm$ hidden-charm pentaquark states $P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{3}{2}^\pm)$ and $P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{1}{2}^\pm)$, respectively,

$$\begin{aligned}
\langle 0 | J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(0) | P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{3}{2}^-)(p) \rangle &= \lambda_P^- U_\mu^-(p, s), \\
\langle 0 | J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(0) | P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{3}{2}^+)(p) \rangle &= \lambda_P^+ i\gamma_5 U_\mu^+(p, s), \\
\langle 0 | J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(0) | P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{1}{2}^+)(p) \rangle &= \tilde{\lambda}_P^+ U^+(p, s) p_\mu, \\
\langle 0 | J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(0) | P_{q_1 q_2 q_3}^{j_L j_H j}(\frac{1}{2}^-)(p) \rangle &= \tilde{\lambda}_P^- i\gamma_5 U^-(p, s) p_\mu,
\end{aligned}$$
(15)

where the $\lambda_P^\pm$ and $\tilde{\lambda}_P^\pm$ are the pole residues or current-hadron coupling constants, the $U_\mu^\pm(p, s)$ and $U^\pm(p, s)$ are the Dirac spinors [22, 23, 24, 25]. The Dirac spinors $U_\mu^\pm(p, s)$ obey the Rarita-Schwinger equation $(\not{p} - M_{P,\pm})U_\mu^\pm(p) = 0$, and the relations $\gamma^\mu U_\mu^\pm(p, s) = 0$, $p^\mu U_\mu^\pm(p, s) = 0$; while the Dirac spinors $U^\pm(p, s)$ obey the Dirac equation $(\not{p} - \widetilde{M}_{P,\pm})U^\pm(p) = 0$.

At the phenomenological side, we can insert a complete set of intermediate hidden-charm pentaquark states with the same quantum numbers as the current operators $J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(x)$ and $i\gamma_5 J_{q_1 q_2 q_3, \mu}^{j_L j_H j}(x)$ into the correlation functions $\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p)$ to obtain the hadronic representation [14, 15]. After isolating the pole terms of the lowest states of the hidden-charm pentaquark states with $J^P = \frac{3}{2}^\pm$ and $J^P = \frac{1}{2}^\pm$ respectively, we obtain the following results:

$$\begin{aligned}
\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p) &= \left(\lambda_P^{-2} \frac{\not{p} + M_{P,-}}{M_{P,-}^2 - p^2} + \lambda_P^{+2} \frac{\not{p} - M_{P,+}}{M_{P,+}^2 - p^2} \right) \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \right) \\
&\quad + \left(\tilde{\lambda}_P^{+2} \frac{\not{p} + \widetilde{M}_{P,+}}{\widetilde{M}_{P,+}^2 - p^2} + \tilde{\lambda}_P^{-2} \frac{\not{p} - \widetilde{M}_{P,-}}{\widetilde{M}_{P,-}^2 - p^2} \right) p_\mu p_\nu + \dots \\
&= \Pi_{q_1 q_2 q_3}^{j_L j_H j}(p^2) \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \right) + \widetilde{\Pi}_{q_1 q_2 q_3}^{j_L j_H j}(p^2) p_\mu p_\nu.
\end{aligned}$$
(16)

In this article, we choose the component $\Pi_{q_1 q_2 q_3}^{j_L j_H j}(p^2)$ associated with the tensor structure $-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2}$ to study the hidden-charm pentaquark states with $J^P = \frac{3}{2}^\pm$, there are no contaminations come from the hidden-charm pentaquark states with $J^P = \frac{1}{2}^\pm$ in the component $\tilde{\Pi}_{q_1 q_2 q_3}^{j_L j_H j}(p^2)$ associated with the tensor structure $p_\mu p_\nu$. It is easy to project out the component $\Pi_{q_1 q_2 q_3}^{j_L j_H j}(p^2)$,

$$\Pi_{q_1 q_2 q_3}^{j_L j_H j}(p^2) = \frac{1}{3} \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \right) \Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p). \quad (17)$$

We can obtain the hadronic spectral densities through dispersion relation,

$$\begin{aligned} \frac{\text{Im} \Pi_{q_1 q_2 q_3}^{j_L j_H j}(s)}{\pi} &= \not{p} \left[\lambda_P^{-2} \delta(s - M_{P,-}^2) + \lambda_P^{+2} \delta(s - M_{P,+}^2) \right] \\ &\quad + \left[M_{P,-} \lambda_P^{-2} \delta(s - M_{P,-}^2) - M_{P,+} \lambda_P^{+2} \delta(s - M_{P,+}^2) \right], \\ &= \not{p} \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 1}(s) + \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 0}(s), \end{aligned} \quad (18)$$

where the subscript index H denotes the hadron side (or phenomenological side). Then we introduce the weight function $\exp(-\frac{s}{T^2})$ to obtain the phenomenological side of the QCD sum rules,

$$\int_{4m_c^2}^{s_0} ds \left[\sqrt{s} \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 1}(s) + \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 0}(s) \right] \exp\left(-\frac{s}{T^2}\right) = 2M_{P,-} \lambda_P^{-2} \exp\left(-\frac{M_{P,-}^2}{T^2}\right), \quad (19)$$

$$\int_{4m_c^2}^{s_0} ds \left[\sqrt{s} \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 1}(s) - \rho_{q_1 q_2 q_3, H}^{j_L j_H j, 0}(s) \right] \exp\left(-\frac{s}{T^2}\right) = 2M_{P,+} \lambda_P^{+2} \exp\left(-\frac{M_{P,+}^2}{T^2}\right), \quad (20)$$

where the T^2 are the Borel parameters and the s_0 are the continuum threshold parameters. In Eqs.(19-20), we separate the contributions of the negative-parity pentaquark states from that of the positive-parity pentaquark states explicitly. They do not contaminate each other, and warrant robust predictions.

Now we briefly outline the operator product expansion for the correlation functions $\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p)$ and use the example $\Pi_{ss\bar{s}, \mu\nu}^{10\frac{1}{2}}(p)$ to illustrate the routine. Firstly, we contract the s and c quark fields in the correlation function $\Pi_{ss\bar{s}, \mu\nu}^{10\frac{1}{2}}(p)$ with Wick theorem,

$$\begin{aligned} \Pi_{ss\bar{s}, \mu\nu}^{10\frac{1}{2}}(p) &= i \varepsilon^{ila} \varepsilon^{ijk} \varepsilon^{lmn} \varepsilon^{i'l'a'} \varepsilon^{i'j'k'} \varepsilon^{l'm'n'} \int d^4x e^{ip \cdot x} C C_{a'a}^T(-x) C \\ &\quad \{ 2Tr [\gamma_\mu S_{kk'}(x) \gamma_\nu C S_{jj'}^T(x) C] Tr [\gamma_5 C_{nn'}(x) \gamma_5 C S_{mm'}^T(x) C] \\ &\quad - 4Tr [\gamma_\mu S_{kk'}(x) \gamma_\nu C S_{mj'}^T(x) C \gamma_5 C_{nn'}(x) \gamma_5 C S_{jm'}^T(x) C] \}, \end{aligned} \quad (21)$$

where the $S_{ij}(x)$ and $C_{ij}(x)$ are the full s and c quark propagators, respectively,

$$\begin{aligned} S_{ij}(x) &= \frac{i\delta_{ij} \not{x}}{2\pi^2 x^4} - \frac{\delta_{ij} m_s}{4\pi^2 x^2} - \frac{\delta_{ij} \langle \bar{s}s \rangle}{12} + \frac{i\delta_{ij} \not{x} m_s \langle \bar{s}s \rangle}{48} - \frac{\delta_{ij} x^2 \langle \bar{s} g_s \sigma G s \rangle}{192} + \frac{i\delta_{ij} x^2 \not{x} m_s \langle \bar{s} g_s \sigma G s \rangle}{1152} \\ &\quad - \frac{ig_s G_{\alpha\beta}^a t_{ij}^a (\not{x} \sigma^{\alpha\beta} + \sigma^{\alpha\beta} \not{x})}{32\pi^2 x^2} - \frac{1}{8} \langle \bar{s}_j \sigma^{\mu\nu} s_i \rangle \sigma_{\mu\nu} + \dots, \end{aligned} \quad (22)$$

$$\begin{aligned} C_{ij}(x) &= \frac{i}{(2\pi)^4} \int d^4k e^{-ik \cdot x} \left\{ \frac{\delta_{ij}}{\not{k} - m_c} - \frac{g_s G_{\alpha\beta}^n t_{ij}^n \sigma^{\alpha\beta} (\not{k} + m_c) + (\not{k} + m_c) \sigma^{\alpha\beta}}{4(k^2 - m_c^2)^2} \right. \\ &\quad \left. - \frac{g_s^2 (t^a t^b)_{ij} G_{\alpha\beta}^a G_{\mu\nu}^b (f^{\alpha\beta\mu\nu} + f^{\alpha\mu\beta\nu} + f^{\alpha\mu\nu\beta})}{4(k^2 - m_c^2)^5} + \dots \right\}, \\ f^{\alpha\beta\mu\nu} &= (\not{k} + m_c) \gamma^\alpha (\not{k} + m_c) \gamma^\beta (\not{k} + m_c) \gamma^\mu (\not{k} + m_c) \gamma^\nu (\not{k} + m_c), \end{aligned} \quad (23)$$

and $t^n = \frac{\lambda^n}{2}$, the λ^n is the Gell-Mann matrix [15], then we compute all the integrals in the coordinate space and in the momentum space to obtain the correlation function $\Pi_{ss,\mu\nu}^{10\frac{1}{2}}(p)$ at the quark level. The calculations are straightforward but tedious. All the correlation functions $\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p)$ at the QCD side can be written as

$$\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p) = \Pi_{q_1 q_2 q_3, QCD}^{j_L j_H j}(p^2) \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \right) + \dots, \quad (24)$$

where we add the notation QCD to denote the QCD side. Once the analytical expressions of the correlation functions $\Pi_{q_1 q_2 q_3, \mu\nu}^{j_L j_H j}(p)$ are gotten, we can obtain the QCD spectral densities $\rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s)$ and $\tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)$ through dispersion relation,

$$\frac{\text{Im} \Pi_{q_1 q_2 q_3, QCD}^{j_L j_H j}(s)}{\pi} = \not{p} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) + m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s). \quad (25)$$

The lengthy expressions of the QCD spectral densities $\rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s)$ and $\tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)$ are given explicitly in the Appendix. We carry out the operator product expansion to the vacuum condensates up to dimension 10, and assume vacuum saturation for the higher dimension vacuum condensates. The gluon condensates are accompanied with large numerical denominators, their contributions to the total QCD spectral densities are less (or much less) than the contributions of the dimension 10 vacuum condensates D_{10} [9, 10], see Eq.(26), so we neglect the vacuum condensates involving the gluon condensate $\langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q}q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{s}s \rangle \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q}q \rangle^2 \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q}q \rangle \langle \bar{s}s \rangle \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{s}s \rangle^2 \langle \frac{\alpha_s GG}{\pi} \rangle$, the predictive ability will not be impaired. In calculations, we take into account the contributions of the terms D_0 , D_3 , D_5 , D_6 , D_8 , D_9 and D_{10} , where

$$\begin{aligned} D_0 &= \text{perturbative terms}, \\ D_3 &\propto \langle \bar{q}q \rangle, \langle \bar{s}s \rangle, \\ D_5 &\propto \langle \bar{q}g_s \sigma G q \rangle, \langle \bar{s}g_s \sigma G s \rangle, \\ D_6 &\propto \langle \bar{q}q \rangle^2, \langle \bar{q}q \rangle \langle \bar{s}s \rangle, \langle \bar{s}s \rangle^2, \\ D_8 &\propto \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma G q \rangle, \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma G q \rangle, \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma G s \rangle, \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma G s \rangle, \\ D_9 &\propto \langle \bar{q}q \rangle^3, \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2, \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle, \langle \bar{s}s \rangle^3, \\ D_{10} &\propto \langle \bar{q}g_s \sigma G q \rangle^2, \langle \bar{q}g_s \sigma G q \rangle \langle \bar{s}g_s \sigma G s \rangle, \langle \bar{s}g_s \sigma G s \rangle^2. \end{aligned} \quad (26)$$

Now we can take the quark-hadron duality below the continuum thresholds s_0 and obtain the following QCD sum rules:

$$2M_{P,-} \lambda_P^{-2} \exp\left(-\frac{M_{P,-}^2}{T^2}\right) = \int_{4m_c^2}^{s_0} ds [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) + m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right), \quad (27)$$

$$2M_{P,+} \lambda_P^{+2} \exp\left(-\frac{M_{P,+}^2}{T^2}\right) = \int_{4m_c^2}^{s_0} ds [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) - m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right). \quad (28)$$

We differentiate Eqs.(27-28) with respect to $\frac{1}{T^2}$, then eliminate the pole residues $\lambda_P^\pm$ and obtain the QCD sum rules for the masses of the hidden-charm pentaquark states $M_{P,\mp}^2$,

$$M_{P,-}^2 = \frac{\int_{4m_c^2}^{s_0} ds \frac{d}{d(-1/T^2)} [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) + m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right)}{\int_{4m_c^2}^{s_0} ds [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) + m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right)}, \quad (29)$$

$$M_{P,+}^2 = \frac{\int_{4m_c^2}^{s_0} ds \frac{d}{d(-1/T^2)} [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) - m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right)}{\int_{4m_c^2}^{s_0} ds [\sqrt{s} \rho_{q_1 q_2 q_3}^{j_L j_H j, 1}(s) - m_c \tilde{\rho}_{q_1 q_2 q_3}^{j_L j_H j, 0}(s)] \exp\left(-\frac{s}{T^2}\right)}. \quad (30)$$

Parameters	Values
$\langle \bar{q}q \rangle (1\text{GeV})$	$-(0.24 \pm 0.01 \text{ GeV})^3$ [14, 15, 16]
$\langle \bar{s}s \rangle (1\text{GeV})$	$(0.8 \pm 0.1) \langle \bar{q}q \rangle (1\text{GeV})$ [14, 15, 16]
$\langle \bar{q}g_s\sigma Gq \rangle (1\text{GeV})$	$m_0^2 \langle \bar{q}q \rangle (1\text{GeV})$ [14, 15, 16]
$\langle \bar{s}g_s\sigma Gs \rangle (1\text{GeV})$	$m_0^2 \langle \bar{s}s \rangle (1\text{GeV})$ [14, 15, 16]
$m_0^2 (1\text{GeV})$	$(0.8 \pm 0.1) \text{ GeV}^2$ [14, 15, 16]
$m_c(m_c)$	$(1.275 \pm 0.025) \text{ GeV}$ [26]
$m_s(2\text{GeV})$	$(0.095 \pm 0.005) \text{ GeV}$ [26]

Table 1: The input parameters in the QCD sum rules, the values in the bracket denote the energy scales $\mu = 1 \text{ GeV}$, 2 GeV and m_c , respectively.

We obtain the masses through fractions, see Eqs.(29-30), the contributions involving the gluon condensate in the numerator and denominator are canceled out with each other, the main effects of the gluon condensates can be safely absorbed into the pole residues $\lambda_P^\pm$ and the remaining tiny effects on the masses can be safely neglected.

3 Numerical results and discussions

The input parameters are shown explicitly in Table 1. The quark condensates, mixed quark condensates and $\overline{MS}$ masses evolve with the renormalization group equation, we take into account the energy-scale dependence according to the following equations,

$$\begin{aligned}
\langle \bar{q}q \rangle(\mu) &= \langle \bar{q}q \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{4}{9}}, \\
\langle \bar{s}s \rangle(\mu) &= \langle \bar{s}s \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{4}{9}}, \\
\langle \bar{q}g_s\sigma Gq \rangle(\mu) &= \langle \bar{q}g_s\sigma Gq \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{2}{27}}, \\
\langle \bar{s}g_s\sigma Gs \rangle(\mu) &= \langle \bar{s}g_s\sigma Gs \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{2}{27}}, \\
m_c(\mu) &= m_c(m_c) \left[\frac{\alpha_s(\mu)}{\alpha_s(m_c)} \right]^{\frac{12}{25}}, \\
m_s(\mu) &= m_s(2\text{GeV}) \left[\frac{\alpha_s(\mu)}{\alpha_s(2\text{GeV})} \right]^{\frac{4}{9}}, \\
\alpha_s(\mu) &= \frac{1}{b_0 t} \left[1 - \frac{b_1}{b_0^2} \frac{\log t}{t} + \frac{b_1^2 (\log^2 t - \log t - 1) + b_0 b_2}{b_0^4 t^2} \right], \tag{31}
\end{aligned}$$

where $t = \log \frac{\mu^2}{\Lambda^2}$, $b_0 = \frac{33-2n_f}{12\pi}$, $b_1 = \frac{153-19n_f}{24\pi^2}$, $b_2 = \frac{2857 - \frac{5033}{9}n_f + \frac{325}{27}n_f^2}{128\pi^3}$, $\Lambda = 213 \text{ MeV}$, 296 MeV and 339 MeV for the flavors $n_f = 5, 4$ and 3 , respectively [26]. Furthermore, we set the $m_u = m_d = 0$.

In Refs.[25, 27], we separate the contributions come from the positive parity and negative parity baryon states explicitly, and study the masses and pole residues of the $J^P = \frac{1}{2}^\pm$ and $\frac{3}{2}^\pm$ heavy, doubly-heavy and triply-heavy baryon states with the QCD sum rules systematically. In calculations, we observe that the continuum threshold parameters $\sqrt{s_0} = M_{\text{gr}} + (0.6-0.8) \text{ GeV}$ can reproduce the experimental values of the masses of the observed heavy baryon states [26], where the subscript gr denotes the ground state baryon states. The pentaquark states are another type

baryon states considering the fractional spins, such as $\frac{1}{2}, \frac{3}{2}, \frac{5}{2}$. In Ref.[9], we take the continuum threshold parameters as $\sqrt{s_0} = M_{P_c(4380/4450)} + (0.6 - 0.8) \text{ GeV}$ in the QCD sum rules for the scalar-diquark-axialvector-diquark-antiquark type hidden-charm pentaquark states, which can reproduce the experimental values of the masses $M_{P_c(4380/4450)}$. In the present QCD sum rules, we take the continuum threshold parameters as $\sqrt{s_0} = M_P + (0.6 - 0.8) \text{ GeV}$.

The hidden-charm or hidden-bottom five-quark systems $q_1 q_2 q_3 Q \bar{Q}$ can be described by a double-well potential in the heavy quark limit. The two light quarks q_1 and q_2 combine together to form a light diquark state or correlation $\mathcal{D}_3^i(q_1 q_2)$ in color antitriplet, the heavy antiquark $\bar{Q}$ serves as one static well potential and combines with the light diquark state $\mathcal{D}_3^i(q_1 q_2)$ to form a heavy antitriquark state or correlation $\varepsilon_{ijk} \mathcal{D}_3^i(q_1 q_2) \bar{Q}_3^j$ in color triplet, where the i, j and k are color indexes. On the other hand, the heavy quark Q serves as the other static well potential and combines with the light quark q_3 to form a heavy diquark state or correlation $\mathcal{D}_3^k(q_3 Q)$ in color antitriplet. The heavy diquark state $\mathcal{D}_3^k(q_3 Q)$ and the heavy antitriquark state $\varepsilon_{ijk} \mathcal{D}_3^i(q_1 q_2) \bar{Q}_3^j$ combine together to form a double heavy physical pentaquark state $\varepsilon_{ijk} \mathcal{D}_3^i(q_1 q_2) \bar{Q}_3^j \mathcal{D}_3^k(q_3 Q)$, the two heavy quarks Q and $\bar{Q}$ stabilize the five-quark systems or pentaquark states, just like the double heavy four-quark systems or tetraquark states [21, 28]. In this article, we construct the interpolating currents according to such routines, see Eqs.(2-13).

In the heavy quark limit, the masses of the hidden-charm pentaquark states can be written as

$$M_P = 2m_c + \bar{\Lambda} + \mathcal{O}\left(\frac{1}{m_c}\right) + \dots, \quad (32)$$

where the $\bar{\Lambda}$ is a parameter of the order $\mathcal{O}(m_c^0)$, and independent on the heavy flavor. The m_c can be taken to be the pole mass according to the reparameterization invariance. The fitted value from the mass of the D -mesons in the heavy quark effective theory is about $m_c = 1.3 \text{ GeV}$ [29], which is approximate to the $\overline{MS}$ mass $m_c(m_c) = (1.275 \pm 0.025) \text{ GeV}$ from the Particle Data Group [26]. The $\overline{MS}$ mass $m_c(m_c)$ relates with the pole mass m_c through the relation [26],

$$m_c = m_c(m_c) \left[1 + \frac{4\alpha_s(m_c)}{3\pi} + \dots \right]. \quad (33)$$

Up to corrections of the order $\mathcal{O}(\alpha_s^3)$, the $\overline{MS}$ mass $m_c(m_c) = (1.275 \pm 0.025) \text{ GeV}$ corresponds to the pole mass $m_c = (1.67 \pm 0.07) \text{ GeV}$ [26], which is quite different from the fitted value $m_c = 1.3 \text{ GeV}$ [29]. The heavy quark mass m_Q is just a parameter.

Now we assume that there exists an effective heavy quark mass $\mathbb{M}_c$ or $\mathbb{M}_b$, which is a free parameter, and is larger than the fitted value of the pole mass $m_c = 1.3 \text{ GeV}$ or $m_b = 4.7 \text{ GeV}$ [29] and differs from the $\overline{MS}$ mass $m_c(m_c)$ or $m_b(m_b)$, some effects of the order $\mathcal{O}(m_c^0)$ or $\mathcal{O}(m_b^0)$ are embodied in the $\mathbb{M}_c$ or $\mathbb{M}_b$. Then the double heavy five-quark systems $q_1 q_2 q_3 Q \bar{Q}$ are characterized by the effective heavy quark masses $\mathbb{M}_Q$ and the virtuality $V = \sqrt{M_P^2 - (2\mathbb{M}_Q)^2}$, just like the double heavy four-quark systems $q_1 \bar{q}_2 Q \bar{Q}$ [21].

The QCD sum rules for the hidden-charm or hidden-bottom pentaquark states $q_1 q_2 q_3 Q \bar{Q}$ have three typical energy scales μ^2, T^2, V^2 , we can set the energy scales to be $\mu^2 = V^2 = \mathcal{O}(T^2)$, and obtain an energy scale formula,

$$\mu = \sqrt{M_P^2 - (2\mathbb{M}_Q)^2}, \quad (34)$$

to determine the ideal energy scales of the QCD spectral densities [9, 10, 11]. Once the pentaquark mass M_P is specialized, the effective heavy quark mass $\mathbb{M}_Q$ determines the energy scale μ , which determines the $\overline{MS}$ mass $m_Q(\mu)$ therefore the pentaquark mass M_P based on the QCD sum rules in return. In this article, we take the energy scale formula $\mu = \sqrt{M_P^2 - (2\mathbb{M}_c)^2}$ as a powerful constraint to obey. In Ref.[9], we choose the value $\mathbb{M}_c = 1.8 \text{ GeV}$ determined in the QCD sum rules for the hidden-charm tetraquark states [21] and obtain the ideal energy scales $\mu = 2.5 \text{ GeV}$

and $\mu = 2.6 \text{ GeV}$ for the pentaquark states $P_c(4380)$ and $P_c(4450)$, respectively. The empirical energy scale formula $\mu = \sqrt{M_{X/Y/Z/P}^2 - (2\mathbb{M}_c)^2}$ works well for the hidden-charm tetraquark states and hidden-charm pentaquark states [9, 10, 11, 21].

In the present QCD sum rules, we choose the Borel parameters T^2 and continuum threshold parameters s_0 to obey the following four criteria:

1. Pole dominance at the phenomenological side;
2. Convergence of the operator product expansion;
3. Appearance of the Borel platforms;
4. Satisfying the energy scale formula.

In the QCD sum rules for the three-quark baryon states $qq'Q$, qQQ' , $QQ'Q''$ [25, 27], the predicted masses increase slowly with the increase of the Borel parameters, the Borel platforms do not appear at the minimum values and are not very flat, we determine the Borel windows by the criteria 1 and 2, the two necessary constraints warrant that the extracted masses are reliable. The pentaquark states are another type baryon states considering the fractional spins, so their Borel platforms maybe not appear at the minimum values of the predicted masses. In this article, we choose small Borel windows $T_{max}^2 - T_{min}^2 = 0.4 \text{ GeV}^2$, just like in the case of the hidden-charm tetraquark states [21] and hidden-charm pentaquark states [9, 10, 11], and obtain the Borel platforms by requiring the uncertainties $\frac{\delta M_P}{M_P}$ induced by the Borel parameters are about 1%, the criterion 3 is modified slightly and satisfied automatically.

We search for the ideal Borel parameters T^2 and continuum threshold parameters s_0 according to the four criteria 1, 2, 3 and 4 using try and error. The resulting Borel parameters or Borel windows T^2 , continuum threshold parameters s_0 , ideal energy scales of the QCD spectral densities, pole contributions of the ground state pentaquark states, and contributions of the vacuum condensates of dimension 9 and 10 in the operator product expansion are shown explicitly in Table 2. From the table, we can see that the first two criteria of the QCD sum rules are satisfied, so we expect to make reasonable predictions.

We take into account all uncertainties of the input parameters, and obtain the masses and pole residues of the $J^P = \frac{3}{2}^\pm$ hidden-charm pentaquark states, which are shown explicitly in Table 3. From Table 2 and Table 3, we can see that the criterion 4 is also satisfied. Now the four criteria of the QCD sum rules are all satisfied, and we expect to make reliable predictions. The present predictions of the hidden-charm pentaquark masses can be confronted to the experimental data in the future.

In the isospin limit,

$$\begin{aligned}
M_P &= 4.39 \pm 0.13 \text{ GeV} \text{ for } P_{uuu}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{udd}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{ddd}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right), \\
M_P &= 4.39 \pm 0.14 \text{ GeV} \text{ for } P_{uuu}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{udd}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right), P_{ddd}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right), \\
M_P &= 4.39 \pm 0.14 \text{ GeV} \text{ for } P_{uuu}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right), P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right), P_{udd}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right), P_{ddd}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right), \quad (35)
\end{aligned}$$

which are all compatible with the experimental value $M_{P_c(4380)} = 4380 \pm 8 \pm 29 \text{ MeV}$ from the LHCb collaboration [1]. The $P_c(4380)$ can be assigned to be the pentaquark state $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ or $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$, or the $P_c(4380)$ has some pentaquark components $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$. From Eq.(35), we can see that the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$ have almost degenerate masses. All the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$ have the light axialvector diquark state $\varepsilon^{ijk}u_j^T C\gamma_\mu u_k$ or $\varepsilon^{ijk}u_j^T C\gamma_\mu d_k$, moreover, the heavy-light scalar and axialvector diquark states have almost degenerate masses, so

the axialvector-diquark-scalar-diquark-antiquark type hidden-charm pentaquark states and the axialvector-diquark-axialvector-diquark-antiquark type hidden-charm pentaquark states maybe have degenerate masses. The mass alone cannot identify the $P_c(4380)$ unambiguously, more experimental data on its productions and decays are still needed. On the theoretical side, we can study the two-body strong decays $P_c(4380) \rightarrow J/\psi p$ with the three-point QCD sum rules or the light-cone QCD sum rules, which maybe give additional support for such assignment.

The Λ_b^0 can be well interpolated by the current $J(x) = \varepsilon^{ijk} u_i^T(x) C \gamma_5 d_j(x) b_k(x)$ [27], the u and d quarks in the Λ_b^0 form a scalar diquark $[ud]$ (or $\varepsilon^{ijk} u_i^T C \gamma_5 d_j$) in color antitriplet, the decays $\Lambda_b^0 \rightarrow J/\psi p K^-$ take place through the following mechanism,

$$\Lambda_b^0([ud]b) \rightarrow [ud]c\bar{c}s \rightarrow [ud]c\bar{c}u\bar{u}s \rightarrow P_c^+([ud][uc]\bar{c})K^-(\bar{u}s) \rightarrow J/\psi p K^-, \quad (36)$$

at the quark level. In the decays $P_c^+([ud][uc]\bar{c}) \rightarrow J/\psi p$, the scalar diquark $[ud]$ survives in the decay chains, the decays are greatly facilitated and can take place easily. On the other hand, if there exists a light axialvector diquark $\{ud\}$ (or $\varepsilon^{ijk} u_i^T C \gamma_\mu d_j$), which has to dissolve to form a scalar diquark $[ud]$, the decays are not facilitated and cannot take place easily, but the decays are not forbidden. It is also possible to assign the $P_c(4380)$ to be the axialvector-diquark-scalar-diquark-antiquark type pentaquark state $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$ or the axialvector-diquark-axialvector-diquark-antiquark type pentaquark states $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$.

In Fig.1, we plot the masses of the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^+\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^+\right)$, $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^+\right)$ with variations of the Borel parameters T^2 as an example. In this article, we take the isospin limit, the masses of the pentaquark states $P_{uud}^{j_L j_H j}\left(\frac{3}{2}^-\right)$ and $P_{uuu}^{j_L j_H j}\left(\frac{3}{2}^-\right)$ degenerate. From the figure, we can see that the platforms in the Borel windows shown in Table 2 appear at the minimum values for the axialvector-diquark-axialvector-diquark-antiquark type pentaquark states with positive parity, and the Borel platforms are very flat; other platforms in the Borel windows shown Table 2 are not as flat, the uncertainties $\frac{\delta M_P}{M_P}$ induced by the Borel parameters in the Borel windows are about 1%. In Fig.1, we plot the masses with variations of the Borel parameters at a large interval $T^2 = (2.6 - 4.6) \text{ GeV}^2$, which is much larger than the interval of the Borel windows shown in Table 2. Compared to the values shown in Table 3, the masses shown in Fig.1 have larger uncertainties due to the larger interval of the Borel parameters. Furthermore, from the values in Table 3, we can see that the mass gaps between the pentaquark states with opposite parity $P_{q_1 q_2 q_3}^{11\frac{1}{2}}\left(\frac{3}{2}^+\right) - P_{q_1 q_2 q_3}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{q_1 q_2 q_3}^{11\frac{3}{2}}\left(\frac{3}{2}^+\right) - P_{q_1 q_2 q_3}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$ are about 0.5 GeV and 0.7 GeV respectively, which is much larger than the mass gap 0.1 GeV between the pentaquark states with opposite parity $P_{q_1 q_2 q_3}^{10\frac{1}{2}}\left(\frac{3}{2}^+\right) - P_{q_1 q_2 q_3}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$. An additional P-wave costs about 0.5 GeV for the conventional baryon states, the mass spectra of the pentaquark states have new feature.

4 Conclusion

In this article, we construct the axialvector-diquark-scalar-diquark-antiquark type and the axialvector-diquark-axialvector-diquark-antiquark type currents to interpolate the $J^P = \frac{3}{2}^\pm$ hidden-charm pentaquark states, study the masses and pole residues with the QCD sum rules systematically by calculating the contributions of the vacuum condensates up to dimension 10 in the operator product expansion. We obtain the masses of the hidden-charm pentaquark states with the strangeness $S = 0, -1, -2, -3$, respectively, which can be confronted to the experimental data in the future. As a byproduct, the predicted pole residues can be taken as basic input parameters in studying the strong decays of the hidden-charm pentaquark states with the three-point QCD sum rules or the light-cone QCD sum rules. In calculations, we use the empirical energy

	$T^2(\text{GeV}^2)$	$\sqrt{s_0}(\text{GeV})$	$\mu(\text{GeV})$	pole	D_9	D_{10}
$P_{uuu}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.3 – 3.7	5.1 ± 0.1	2.5	(41 – 62)%	(11 – 15)%	(3 – 4)%
$P_{uus}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.5 – 3.9	5.2 ± 0.1	2.7	(40 – 60)%	(7 – 9)%	(2 – 3)%
$P_{uss}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.6 – 4.0	5.3 ± 0.1	2.8	(41 – 61)%	(5 – 6)%	(1 – 2)%
$P_{sss}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.8 – 4.2	5.4 ± 0.1	3.0	(40 – 59)%	(3 – 4)%	$\sim 1\%$
$P_{uuu}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.2 – 3.6	5.1 ± 0.1	2.5	(42 – 64)%	(12 – 17)%	(3 – 5)%
$P_{uus}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.4 – 3.8	5.2 ± 0.1	2.7	(41 – 61)%	(7 – 10)%	(2 – 3)%
$P_{uss}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.6 – 4.0	5.3 ± 0.1	2.9	(40 – 59)%	(4 – 6)%	(1 – 2)%
$P_{sss}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	3.7 – 4.1	5.4 ± 0.1	3.0	(41 – 60)%	(3 – 4)%	(1 – 2)%
$P_{uuu}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	3.2 – 3.6	5.1 ± 0.1	2.5	(41 – 63)%	(12 – 17)%	(2 – 3)%
$P_{uus}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	3.4 – 3.8	5.2 ± 0.1	2.7	(40 – 60)%	(7 – 10)%	(1 – 2)%
$P_{uss}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	3.5 – 3.9	5.3 ± 0.1	2.9	(41 – 61)%	(5 – 7)%	$\sim 1\%$
$P_{sss}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	3.7 – 4.1	5.4 ± 0.1	3.0	(40 – 59)%	(3 – 4)%	$\sim 1\%$
$P_{uuu}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.2 – 3.6	5.2 ± 0.1	2.7	(41 – 62)%	(13 – 18)%	(3 – 5)%
$P_{uus}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.3 – 3.7	5.3 ± 0.1	2.9	(42 – 63)%	(9 – 12)%	(2 – 4)%
$P_{uss}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.5 – 3.9	5.4 ± 0.1	3.0	(40 – 60)%	(6 – 8)%	$\sim 2\%$
$P_{sss}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.6 – 4.0	5.5 ± 0.1	3.2	(41 – 61)%	(4 – 5)%	(1 – 2)%
$P_{uuu}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.4 – 3.8	5.6 ± 0.1	3.3	(50 – 69)%	–(4 – 6)%	(3 – 4)%
$P_{uus}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.6 – 4.0	5.7 ± 0.1	3.5	(48 – 68)%	–(2 – 4)%	(2 – 3)%
$P_{uss}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	3.8 – 4.2	5.8 ± 0.1	3.6	(47 – 66)%	–(1 – 2)%	(1 – 2)%
$P_{sss}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.0 – 4.4	5.9 ± 0.1	3.7	(46 – 64)%	–(1 – 2)%	$\sim 1\%$
$P_{uuu}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	3.3 – 3.7	5.8 ± 0.1	3.7	(60 – 78)%	–(3 – 6)%	$\sim -1\%$
$P_{uus}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	3.5 – 3.9	5.9 ± 0.1	3.8	(59 – 77)%	–(2 – 3)%	$\sim -1\%$
$P_{uss}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	3.7 – 4.1	6.0 ± 0.1	3.9	(58 – 75)%	–(1 – 2)%	$\sim -0\%$
$P_{sss}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	3.9 – 4.3	6.1 ± 0.1	4.0	(57 – 74)%	–1%	$\sim -0\%$

Table 2: The Borel parameters (Borel windows), continuum threshold parameters, ideal energy scales, pole contributions, contributions of the vacuum condensates of dimension 9 and dimension 10 for the hidden-charm pentaquark states.

	$M_P(\text{GeV})$	$\lambda_P(10^{-3}\text{GeV}^6)$
$P_{uuu}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.39 ± 0.13	2.25 ± 0.40
$P_{uus}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.51 ± 0.12	2.75 ± 0.45
$P_{uss}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.60 ± 0.11	3.19 ± 0.50
$P_{sss}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.70 ± 0.11	3.78 ± 0.57
$P_{uuu}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.39 ± 0.14	3.75 ± 0.68
$P_{uus}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.51 ± 0.12	4.64 ± 0.77
$P_{uss}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.62 ± 0.11	5.60 ± 0.88
$P_{sss}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$	4.71 ± 0.11	6.47 ± 1.00
$P_{uuu}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	4.39 ± 0.14	3.74 ± 0.70
$P_{uus}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	4.52 ± 0.12	4.64 ± 0.79
$P_{uss}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	4.61 ± 0.12	5.52 ± 0.90
$P_{sss}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$	4.72 ± 0.11	6.52 ± 1.02
$P_{uuu}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.51 ± 0.13	0.99 ± 0.19
$P_{uus}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.60 ± 0.12	1.21 ± 0.22
$P_{uss}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.72 ± 0.11	1.45 ± 0.25
$P_{sss}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.81 ± 0.12	1.71 ± 0.29
$P_{uuu}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.91 ± 0.09	5.04 ± 0.68
$P_{uus}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	4.99 ± 0.09	5.83 ± 0.82
$P_{uss}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	5.08 ± 0.09	6.72 ± 0.98
$P_{sss}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$	5.18 ± 0.09	7.73 ± 1.10
$P_{uuu}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	5.14 ± 0.08	7.86 ± 0.91
$P_{uus}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	5.21 ± 0.08	8.79 ± 1.07
$P_{uss}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	5.28 ± 0.08	9.91 ± 1.25
$P_{sss}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$	5.36 ± 0.08	11.20 ± 1.42

Table 3: The masses and pole residues of the hidden-charm pentaquark states.




Figure 1: The masses of the pentaquark states with variations of the Borel parameters T^2 , where the A, B, C, D, E and F denote the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$, $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^{-}\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^{+}\right)$, $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^{-}\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^{+}\right)$ respectively, the horizontal lines denote the experimental value of the mass of the $P_c(4380)$.

scale formula $\mu = \sqrt{M_{X/Y/Z/P}^2 - (2\mathbb{M}_c)^2}$ with the effective heavy quark mass $\mathbb{M}_c$ to determine the ideal energy scales of the QCD spectral densities, which works well in studying the hidden-charm tetraquark states and hidden-charm pentaquark states. The predicted masses of the pentaquark states $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$ are compatible with the experimental value $M_{P_c(4380)} = 4380 \pm 8 \pm 29 \text{ MeV}$ from the LHCb collaboration. We can draw the conclusion tentatively that the $P_c(4380)$ can be assigned to be the pentaquark state $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ or $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$, or the $P_c(4380)$ has some pentaquark components $P_{uud}^{10\frac{1}{2}}\left(\frac{3}{2}^-\right)$, $P_{uud}^{11\frac{1}{2}}\left(\frac{3}{2}^-\right)$ and $P_{uud}^{11\frac{3}{2}}\left(\frac{3}{2}^-\right)$. More experimental data on its productions and decays are still needed to identify the $P_c(4380)$ unambiguously.

Appendix

The QCD spectral densities $\rho_{ss\bar{s}}^{11\frac{3}{2},1}(s)$, $\tilde{\rho}_{ss\bar{s}}^{11\frac{3}{2},0}(s)$, $\rho_{us\bar{s}}^{11\frac{3}{2},1}(s)$, $\tilde{\rho}_{us\bar{s}}^{11\frac{3}{2},0}(s)$, $\rho_{uus\bar{s}}^{11\frac{3}{2},1}(s)$, $\tilde{\rho}_{uus\bar{s}}^{11\frac{3}{2},0}(s)$, $\rho_{uuu\bar{s}}^{11\frac{3}{2},1}(s)$, $\tilde{\rho}_{uuu\bar{s}}^{11\frac{3}{2},0}(s)$, $\rho_{ss\bar{s}}^{11\frac{1}{2},1}(s)$, $\tilde{\rho}_{ss\bar{s}}^{11\frac{1}{2},0}(s)$, $\rho_{us\bar{s}}^{11\frac{1}{2},1}(s)$, $\tilde{\rho}_{us\bar{s}}^{11\frac{1}{2},0}(s)$, $\rho_{uus\bar{s}}^{11\frac{1}{2},1}(s)$, $\tilde{\rho}_{uus\bar{s}}^{11\frac{1}{2},0}(s)$, $\rho_{uuu\bar{s}}^{11\frac{1}{2},1}(s)$, $\tilde{\rho}_{uuu\bar{s}}^{11\frac{1}{2},0}(s)$, $\rho_{ss\bar{s}}^{10\frac{1}{2},1}(s)$, $\tilde{\rho}_{ss\bar{s}}^{10\frac{1}{2},0}(s)$, $\rho_{us\bar{s}}^{10\frac{1}{2},1}(s)$, $\tilde{\rho}_{us\bar{s}}^{10\frac{1}{2},0}(s)$, $\rho_{uus\bar{s}}^{10\frac{1}{2},1}(s)$, $\tilde{\rho}_{uus\bar{s}}^{10\frac{1}{2},0}(s)$, $\rho_{uuu\bar{s}}^{10\frac{1}{2},1}(s)$ and $\tilde{\rho}_{uuu\bar{s}}^{10\frac{1}{2},0}(s)$ of the hidden-charm pentaquark states,

$$\begin{aligned}
\rho_{ss\bar{s}}^{11\frac{3}{2},1}(s) = & \frac{1}{122880\pi^8} \int dydz yz(1-y-z)^4 (s - \bar{m}_c^2)^4 (7s - 2\bar{m}_c^2) \\
& + \frac{1}{614400\pi^8} \int dydz yz(1-y-z)^5 (s - \bar{m}_c^2)^3 (63s^2 - 51s\bar{m}_c^2 + 8\bar{m}_c^4) \\
& - \frac{m_c \langle \bar{s}s \rangle}{768\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& - \frac{m_c \langle \bar{s}s \rangle}{768\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& + \frac{11m_c \langle \bar{s}g_s \sigma G s \rangle}{4096\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2)^2 \\
& + \frac{11m_c \langle \bar{s}g_s \sigma G s \rangle}{8192\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle^2}{24\pi^4} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma G s \rangle}{768\pi^4} \int dydz yz (3s - 2\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma G s \rangle}{1536\pi^4} \int dydz (y+z)(1-y-z) (5s - 4\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s}s \rangle^3}{36\pi^2} \int dy \\
& - \frac{13\langle \bar{s}g_s \sigma G s \rangle^2}{13824\pi^4} \int dydz (y+z) \left[1 + \frac{s}{4} \delta(s - \bar{m}_c^2) \right] \\
& + \frac{11\langle \bar{s}g_s \sigma G s \rangle^2}{1536\pi^4} \int dy y(1-y) \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right] \\
& + \frac{m_s m_c}{12288\pi^8} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s m_c}{49152\pi^8} \int dydz (y+z)(1-y-z)^4 (s-\bar{m}_c^2)^3 (7s-3\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{768\pi^6} \int dydz yz(1-y-z)^2 (s-\bar{m}_c^2)^2 (5s-2\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}s \rangle}{768\pi^6} \int dydz yz(1-y-z)^3 (s-\bar{m}_c^2) (35s^2-37s\bar{m}_c^2+8\bar{m}_c^4) \\
& + \frac{11m_s \langle \bar{s}g_s \sigma Gs \rangle}{1024\pi^6} \int dydz yz(1-y-z) (s-\bar{m}_c^2) (2s-\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{256\pi^6} \int dydz yz(1-y-z)^2 (12s^2-15s\bar{m}_c^2+4\bar{m}_c^4) \\
& - \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{4096\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\bar{m}_c^2) (3s-2\bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle^2}{24\pi^4} \int dydz (y+z) (s-\bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}s \rangle^2}{48\pi^4} \int dydz (y+z)(1-y-z) (4s-3\bar{m}_c^2) \\
& + \frac{89m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{3072\pi^4} \int dydz (y+z) \left[1 + \frac{s}{3} \delta(s-\bar{m}_c^2) \right] \\
& + \frac{m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{768\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& - \frac{199m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{9216\pi^4} \int dy \\
& + \frac{m_s \langle \bar{s}s \rangle^3}{18\pi^2} \int dy y(1-y) \left[1 + \frac{s}{2} \delta(s-\tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{3072\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s-\tilde{m}_c^2) \\
& + \frac{17m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{4608\pi^4} \int dy \left(1 + \frac{s}{2T^2} \right) \delta(s-\tilde{m}_c^2) , \tag{37}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{sss}^{11\frac{3}{2},0}(s) &= \frac{1}{122880\pi^8} \int dydz (y+z)(1-y-z)^4 (s-\bar{m}_c^2)^4 (6s-\bar{m}_c^2) \\
& - \frac{1}{122880\pi^8} \int dydz (y+z)(1-y-z)^5 (s-\bar{m}_c^2)^3 (48s^2-31s\bar{m}_c^2+3\bar{m}_c^4) \\
& - \frac{m_c \langle \bar{s}s \rangle}{192\pi^6} \int dydz (1-y-z)^2 (s-\bar{m}_c^2)^3 \\
& + \frac{m_c \langle \bar{s}s \rangle}{1152\pi^6} \int dydz (1-y-z)^3 (s-\bar{m}_c^2)^2 (5s-2\bar{m}_c^2) \\
& + \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{1024\pi^6} \int dydz (1-y-z) (s-\bar{m}_c^2)^2 \\
& - \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{2048\pi^6} \int dydz (1-y-z)^2 (s-\bar{m}_c^2) (2s-\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle^2}{48\pi^4} \int dydz (y+z)(1-y-z) (s-\bar{m}_c^2) (3s-\bar{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& -\frac{19\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{768\pi^4}\int dydz(y+z)(2s-\overline{m}_c^2) \\
& -\frac{m_c\langle\bar{s}s\rangle^3}{9\pi^2}\int dy \\
& +\frac{11\langle\bar{s}g_s\sigma Gs\rangle^2}{3072\pi^4}\int dy[1+s\delta(s-\tilde{m}_c^2)] \\
& +\frac{m_sm_c}{3072\pi^8}\int dydz(1-y-z)^3(s-\overline{m}_c^2)^4 \\
& -\frac{m_sm_c}{12288\pi^8}\int dydz(1-y-z)^4(s-\overline{m}_c^2)^3(3s-\overline{m}_c^2) \\
& -\frac{m_s\langle\bar{s}s\rangle}{768\pi^6}\int dydz(y+z)(1-y-z)^2(s-\overline{m}_c^2)^2(4s-\overline{m}_c^2) \\
& -\frac{m_s\langle\bar{s}s\rangle}{512\pi^6}\int dydz(y+z)(1-y-z)^3(s-\overline{m}_c^2)(8s^2-7s\overline{m}_c^2+\overline{m}_c^4) \\
& +\frac{11m_s\langle\bar{s}g_s\sigma Gs\rangle}{2048\pi^6}\int dydz(y+z)(1-y-z)(s-\overline{m}_c^2)(3s-\overline{m}_c^2) \\
& +\frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{1024\pi^6}\int dydz(y+z)(1-y-z)^2(15s^2-16s\overline{m}_c^2+3\overline{m}_c^4) \\
& +\frac{m_sm_c\langle\bar{s}s\rangle^2}{6\pi^4}\int dydz(s-\overline{m}_c^2) \\
& +\frac{m_sm_c\langle\bar{s}s\rangle^2}{24\pi^4}\int dydz(1-y-z)(3s-2\overline{m}_c^2) \\
& -\frac{89m_sm_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{2304\pi^4}\int dydz\left[1+\frac{s}{2}\delta(s-\overline{m}_c^2)\right] \\
& -\frac{199m_sm_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{2304\pi^4}\int dy \\
& +\frac{m_s\langle\bar{s}s\rangle^3}{36\pi^2}\int dy[1+s\delta(s-\tilde{m}_c^2)] \\
& +\frac{125m_sm_c\langle\bar{s}g_s\sigma Gs\rangle^2}{9216\pi^4}\int dy\left(1+\frac{s}{T^2}\right)\delta(s-\tilde{m}_c^2), \tag{38}
\end{aligned}$$

$$\begin{aligned}
\rho_{uss}^{11\frac{3}{2},1}(s) &= \frac{1}{122880\pi^8}\int dydz yz(1-y-z)^4(s-\overline{m}_c^2)^4(7s-2\overline{m}_c^2) \\
& +\frac{1}{614400\pi^8}\int dydz yz(1-y-z)^5(s-\overline{m}_c^2)^3(63s^2-51s\overline{m}_c^2+8\overline{m}_c^4) \\
& -\frac{m_c[\langle\bar{q}q\rangle+2\langle\bar{s}s\rangle]}{2304\pi^6}\int dydz(y+z)(1-y-z)^2(s-\overline{m}_c^2)^3 \\
& -\frac{m_c[\langle\bar{q}q\rangle+2\langle\bar{s}s\rangle]}{2304\pi^6}\int dydz(y+z)(1-y-z)^3(s-\overline{m}_c^2)^2(2s-\overline{m}_c^2) \\
& +\frac{11m_c[\langle\bar{q}g_s\sigma Gq\rangle+2\langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6}\int dydz(y+z)(1-y-z)(s-\overline{m}_c^2)^2 \\
& +\frac{11m_c[\langle\bar{q}g_s\sigma Gq\rangle+2\langle\bar{s}g_s\sigma Gs\rangle]}{24576\pi^6}\int dydz(y+z)(1-y-z)^2(s-\overline{m}_c^2)(5s-3\overline{m}_c^2) \\
& +\frac{\langle\bar{s}s\rangle[2\langle\bar{q}q\rangle+\langle\bar{s}s\rangle]}{72\pi^4}\int dydz yz(1-y-z)(s-\overline{m}_c^2)(2s-\overline{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{2304\pi^4} \int dydz yz (3s - 2\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{4608\pi^4} \int dydz (y+z)(1-y-z) (5s - 4\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{36\pi^2} \int dy \\
& - \frac{13 \langle \bar{s}g_s \sigma Gs \rangle [2 \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{41472\pi^4} \int dydz (y+z) \left[1 + \frac{s}{4} \delta(s - \bar{m}_c^2) \right] \\
& + \frac{11 \langle \bar{s}g_s \sigma Gs \rangle [2 \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{4608\pi^4} \int dy y(1-y) \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c}{18432\pi^8} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& + \frac{m_s m_c}{73728\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^3 (7s - 3\bar{m}_c^2) \\
& - \frac{m_s [2 \langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{1152\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}s \rangle}{1152\pi^6} \int dydz yz(1-y-z)^3 (s - \bar{m}_c^2) (35s^2 - 37s\bar{m}_c^2 + 8\bar{m}_c^4) \\
& - \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{384\pi^6} \int dydz yz(1-y-z)^2 (12s^2 - 15s\bar{m}_c^2 + 4\bar{m}_c^4) \\
& - \frac{m_s [\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (3s - 2\bar{m}_c^2) \\
& + \frac{m_s [19 \langle \bar{q}g_s \sigma Gq \rangle + 3 \langle \bar{s}g_s \sigma Gs \rangle]}{3072\pi^6} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [5 \langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z)(1-y-z) (4s - 3\bar{m}_c^2) \\
& + \frac{m_s m_c [89 \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + 57 \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle + 32 \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{9216\pi^4} \\
& \int dydz (y+z) \left[1 + \frac{s}{3} \delta(s - \bar{m}_c^2) \right] \\
& - \frac{m_s m_c [231 \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 89 \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + 256 \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c [\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{2304\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{27\pi^2} \int dy y(1-y) \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{4608\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle [19 \langle \bar{s}g_s \sigma Gs \rangle - 53 \langle \bar{q}g_s \sigma Gq \rangle]}{13824\pi^4} \int dy \left(1 + \frac{s}{2T^2} \right) \delta(s - \tilde{m}_c^2) , \tag{39}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uss}^{11\frac{3}{2},0}(s) = & \frac{1}{122880\pi^8} \int dydz (y+z)(1-y-z)^4 (s-\bar{m}_c^2)^4 (6s-\bar{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dydz (y+z)(1-y-z)^5 (s-\bar{m}_c^2)^3 (48s^2-31s\bar{m}_c^2+3\bar{m}_c^4) \\
& - \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{576\pi^6} \int dydz (1-y-z)^2 (s-\bar{m}_c^2)^3 \\
& + \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{3456\pi^6} \int dydz (1-y-z)^3 (s-\bar{m}_c^2)^2 (5s-2\bar{m}_c^2) \\
& + \frac{11m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{3072\pi^6} \int dydz (1-y-z) (s-\bar{m}_c^2)^2 \\
& - \frac{11m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{6144\pi^6} \int dydz (1-y-z)^2 (s-\bar{m}_c^2) (2s-\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z)(1-y-z) (s-\bar{m}_c^2) (3s-\bar{m}_c^2) \\
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{2304\pi^4} \int dydz (y+z) (2s-\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{9\pi^2} \int dy \\
& + \frac{11 \langle \bar{s}g_s\sigma Gs \rangle [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{9216\pi^4} \int dy [1+s\delta(s-\tilde{m}_c^2)] \\
& + \frac{m_s m_c}{4608\pi^8} \int dydz (1-y-z)^3 (s-\bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{18432\pi^8} \int dydz (1-y-z)^4 (s-\bar{m}_c^2)^3 (3s-\bar{m}_c^2) \\
& - \frac{m_s [2\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{1152\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\bar{m}_c^2)^2 (4s-\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{768\pi^6} \int dydz (y+z)(1-y-z)^3 (s-\bar{m}_c^2) (8s^2-7s\bar{m}_c^2+\bar{m}_c^4) \\
& + \frac{m_s \langle \bar{s}g_s\sigma Gs \rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^2 (15s^2-16s\bar{m}_c^2+3\bar{m}_c^4) \\
& + \frac{m_s [19\langle \bar{q}g_s\sigma Gq \rangle + 3\langle \bar{s}g_s\sigma Gs \rangle]}{6144\pi^6} \int dydz (y+z)(1-y-z) (s-\bar{m}_c^2) (3s-\bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [5\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{36\pi^4} \int dydz (s-\bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{72\pi^4} \int dydz (1-y-z) (3s-2\bar{m}_c^2) \\
& - \frac{m_s m_c [89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle + 32\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{6912\pi^4} \int dydz \left[1 + \frac{s}{2}\delta(s-\bar{m}_c^2)\right] \\
& - \frac{m_s m_c [231\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle - 89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 256\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{6912\pi^4} \int dy \\
& + \frac{m_s \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{54\pi^2} \int dy [1+s\delta(s-\tilde{m}_c^2)] \\
& - \frac{m_s m_c \langle \bar{s}g_s\sigma Gs \rangle [19\langle \bar{s}g_s\sigma Gs \rangle - 269\langle \bar{q}g_s\sigma Gq \rangle]}{27648\pi^4} \int dy \left(1 + \frac{s}{T^2}\right) \delta(s-\tilde{m}_c^2), \tag{40}
\end{aligned}$$

$$\begin{aligned}
\rho_{uus}^{11\frac{3}{2},1}(s) = & \frac{1}{122880\pi^8} \int dydz yz(1-y-z)^4 (s-\bar{m}_c^2)^4 (7s-2\bar{m}_c^2) \\
& + \frac{1}{614400\pi^8} \int dydz yz(1-y-z)^5 (s-\bar{m}_c^2)^3 (63s^2-51s\bar{m}_c^2+8\bar{m}_c^4) \\
& - \frac{m_c [2\langle\bar{q}q\rangle + \langle\bar{s}s\rangle]}{2304\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\bar{m}_c^2)^3 \\
& - \frac{m_c [2\langle\bar{q}q\rangle + \langle\bar{s}s\rangle]}{2304\pi^6} \int dydz (y+z)(1-y-z)^3 (s-\bar{m}_c^2)^2 (2s-\bar{m}_c^2) \\
& + \frac{11m_c [2\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z) (s-\bar{m}_c^2)^2 \\
& + \frac{11m_c [2\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{24576\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\bar{m}_c^2) (5s-3\bar{m}_c^2) \\
& + \frac{\langle\bar{q}q\rangle [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{72\pi^4} \int dydz yz(1-y-z) (s-\bar{m}_c^2) (2s-\bar{m}_c^2) \\
& - \frac{19 [\langle\bar{q}q\rangle \langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{q}q\rangle \langle\bar{s}g_s\sigma Gs\rangle + \langle\bar{s}s\rangle \langle\bar{q}g_s\sigma Gq\rangle]}{2304\pi^4} \int dydz yz (3s-2\bar{m}_c^2) \\
& + \frac{\langle\bar{q}q\rangle \langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{q}q\rangle \langle\bar{s}g_s\sigma Gs\rangle + \langle\bar{s}s\rangle \langle\bar{q}g_s\sigma Gq\rangle}{4608\pi^4} \int dydz (y+z)(1-y-z) (5s-4\bar{m}_c^2) \\
& - \frac{13\langle\bar{q}g_s\sigma Gq\rangle [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{41472\pi^4} \int dydz (y+z) \left[1 + \frac{s}{4}\delta(s-\bar{m}_c^2)\right] \\
& + \frac{11\langle\bar{q}g_s\sigma Gq\rangle [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{4608\pi^4} \int dy y(1-y) \left[1 + \frac{s}{2}\delta(s-\bar{m}_c^2)\right] \\
& - \frac{m_c \langle\bar{q}q\rangle^2 \langle\bar{s}s\rangle}{36\pi^2} \int dy \\
& + \frac{m_s m_c}{36864\pi^8} \int dydz (y+z)(1-y-z)^3 (s-\bar{m}_c^2)^4 \\
& + \frac{m_s m_c}{147456\pi^8} \int dydz (y+z)(1-y-z)^4 (s-\bar{m}_c^2)^3 (7s-3\bar{m}_c^2) \\
& - \frac{m_s [4\langle\bar{q}q\rangle - 3\langle\bar{s}s\rangle]}{2304\pi^6} \int dydz yz(1-y-z)^2 (s-\bar{m}_c^2)^2 (5s-2\bar{m}_c^2) \\
& + \frac{m_s \langle\bar{s}s\rangle}{2304\pi^6} \int dydz yz(1-y-z)^3 (s-\bar{m}_c^2) (35s^2-37s\bar{m}_c^2+8\bar{m}_c^4) \\
& + \frac{m_s [19\langle\bar{q}g_s\sigma Gq\rangle - 8\langle\bar{s}g_s\sigma Gs\rangle]}{3072\pi^6} \int dydz yz(1-y-z) (s-\bar{m}_c^2) (2s-\bar{m}_c^2) \\
& - \frac{m_s \langle\bar{s}g_s\sigma Gs\rangle}{768\pi^6} \int dydz yz(1-y-z)^2 (12s^2-15s\bar{m}_c^2+4\bar{m}_c^4) \\
& - \frac{m_s \langle\bar{q}g_s\sigma Gq\rangle}{12288\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\bar{m}_c^2) (3s-2\bar{m}_c^2) \\
& + \frac{m_s m_c \langle\bar{q}q\rangle [3\langle\bar{q}q\rangle - \langle\bar{s}s\rangle]}{144\pi^4} \int dydz (y+z) (s-\bar{m}_c^2) \\
& - \frac{m_s m_c \langle\bar{q}q\rangle \langle\bar{s}s\rangle}{144\pi^4} \int dydz (y+z)(1-y-z) (4s-3\bar{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s m_c [32 \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 57 \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{9216\pi^4} \int dy dz (y+z) \left[1 + \frac{s}{3} \delta(s - \bar{m}_c^2) \right] \\
& - \frac{m_s m_c [288 \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle - 57 \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 32 \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle}{2304\pi^4} \int dy dz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{54\pi^2} \int dy y(1-y) \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle^2}{9216\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle [36 \langle \bar{q}g_s \sigma Gq \rangle - 19 \langle \bar{s}g_s \sigma Gs \rangle]}{13824\pi^4} \int dy \left(1 + \frac{s}{2T^2} \right) \delta(s - \tilde{m}_c^2), \tag{41}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uus}^{11\frac{3}{2},0}(s) = & \frac{1}{122880\pi^8} \int dy dz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 (6s - \bar{m}_c^2) \\
& - \frac{1}{122880\pi^8} \int dy dz (y+z)(1-y-z)^5 (s - \bar{m}_c^2)^3 (48s^2 - 31s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& - \frac{m_c [2 \langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{576\pi^6} \int dy dz (1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c [2 \langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{3456\pi^6} \int dy dz (1-y-z)^3 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& + \frac{11m_c [2 \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{3072\pi^6} \int dy dz (1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c [2 \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{6144\pi^6} \int dy dz (1-y-z)^2 (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{\langle \bar{q}q \rangle [\langle \bar{q}q \rangle + 2 \langle \bar{s}s \rangle]}{144\pi^4} \int dy dz (y+z)(1-y-z) (s - \bar{m}_c^2) (3s - \bar{m}_c^2) \\
& - \frac{19 [\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{2304\pi^4} \int dy dz (y+z) (2s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{9\pi^2} \int dy \\
& + \frac{11 \langle \bar{q}g_s \sigma Gq \rangle [\langle \bar{q}g_s \sigma Gq \rangle + 2 \langle \bar{s}g_s \sigma Gs \rangle]}{9216\pi^4} \int dy [1 + s \delta(s - \tilde{m}_c^2)] \\
& + \frac{m_s m_c}{9216\pi^8} \int dy dz (1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{36864\pi^8} \int dy dz (1-y-z)^4 (s - \bar{m}_c^2)^3 (3s - \bar{m}_c^2) \\
& - \frac{m_s [4 \langle \bar{q}q \rangle - 3 \langle \bar{s}s \rangle]}{2304\pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 (4s - \bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{1536\pi^6} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2) (8s^2 - 7s\bar{m}_c^2 + \bar{m}_c^4) \\
& + \frac{m_s [19 \langle \bar{q}g_s \sigma Gq \rangle - 8 \langle \bar{s}g_s \sigma Gs \rangle]}{6144\pi^6} \int dy dz (y+z)(1-y-z) (s - \bar{m}_c^2) (3s - \bar{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{3072 \pi^6} \int dy dz (y+z)(1-y-z)^2 (15s^2 - 16s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& + \frac{m_s m_c \langle \bar{q} q \rangle [3\langle \bar{q} q \rangle - \langle \bar{s} s \rangle]}{36 \pi^4} \int dy dz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle}{72 \pi^4} \int dy dz (1-y-z) (3s - 2\bar{m}_c^2) \\
& - \frac{m_s m_c [32\langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + 57\langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle]}{6912 \pi^4} \int dy dz \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right] \\
& - \frac{m_s m_c [288\langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle - 57\langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle - 32\langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle]}{6912 \pi^4} \int dy \\
& + \frac{m_s \langle \bar{q} q \rangle^2 \langle \bar{s} s \rangle}{108 \pi^2} \int dy [1 + s \delta(s - \tilde{m}_c^2)] \\
& + \frac{m_s m_c \langle \bar{q} g_s \sigma G q \rangle [144\langle \bar{q} g_s \sigma G q \rangle - 19\langle \bar{s} g_s \sigma G s \rangle]}{27648 \pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2), \quad (42)
\end{aligned}$$

$$\rho_{uu\bar{u}}^{11\frac{3}{2},1}(s) = \rho_{ss\bar{s}}^{11\frac{3}{2},1}(s) \mid_{m_s \rightarrow 0, \langle \bar{s} s \rangle \rightarrow \langle \bar{q} q \rangle, \langle \bar{s} g_s \sigma G s \rangle \rightarrow \langle \bar{q} g_s \sigma G q \rangle}, \quad (43)$$

$$\hat{\rho}_{uu\bar{u}}^{11\frac{3}{2},0}(s) = \hat{\rho}_{ss\bar{s}}^{11\frac{3}{2},0}(s) \mid_{m_s \rightarrow 0, \langle \bar{s} s \rangle \rightarrow \langle \bar{q} q \rangle, \langle \bar{s} g_s \sigma G s \rangle \rightarrow \langle \bar{q} g_s \sigma G q \rangle}, \quad (44)$$

$$\begin{aligned}
\rho_{ss\bar{s}}^{11\frac{1}{2},1}(s) &= \frac{1}{61440 \pi^8} \int dy dz yz(1-y-z)^4 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{1}{307200 \pi^8} \int dy dz yz(1-y-z)^5 (s - \bar{m}_c^2)^4 (9s - 4\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s} s \rangle}{768 \pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c \langle \bar{s} s \rangle}{2304 \pi^6} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c \langle \bar{s} g_s \sigma G s \rangle}{4096 \pi^6} \int dy dz (y+z)(1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c \langle \bar{s} g_s \sigma G s \rangle}{8192 \pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s} s \rangle^2}{24 \pi^4} \int dy dz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{19\langle \bar{s} s \rangle \langle \bar{s} g_s \sigma G s \rangle}{384 \pi^4} \int dy dz yz (4s - 3\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s} s \rangle^3}{36 \pi^2} \int dy \\
& + \frac{11\langle \bar{s} g_s \sigma G s \rangle^2}{512 \pi^4} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c}{12288 \pi^8} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{49152 \pi^8} \int dy dz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s \langle \bar{s} s \rangle}{128 \pi^6} \int dy dz yz(1-y-z)^2 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& -\frac{m_s \langle \bar{s}s \rangle}{384\pi^6} \int dydz yz(1-y-z)^3 (s - \bar{m}_c^2)^2 (7s - 4\bar{m}_c^2) \\
& + \frac{11m_s \langle \bar{s}g_s \sigma Gs \rangle}{1024\pi^6} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{128\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2) (3s - 2\bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle^2}{24\pi^4} \int dydz (y+z) (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle^2}{48\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) \\
& - \frac{89m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{9216\pi^4} \int dydz (y+z) \\
& - \frac{199m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{9216\pi^4} \int dy \\
& + \frac{m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{768\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& + \frac{m_s \langle \bar{s}s \rangle^3}{6\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{2304\pi^4} \int dy \left(16 + \frac{7s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{3072\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{6144\pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) , \tag{45}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{ss\bar{s}}^{11\frac{1}{2},0}(s) &= \frac{1}{245760\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 (7s - 2\bar{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s}s \rangle}{192\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c \langle \bar{s}s \rangle}{576\pi^6} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{1024\pi^6} \int dydz (1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{2048\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle^2}{48\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{1536\pi^4} \int dydz (y+z) (3s - 2\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{768\pi^4} \int dydz (1-y-z) s \\
& - \frac{m_c \langle \bar{s}s \rangle^3}{9\pi^2} \int dy
\end{aligned}$$

$$\begin{aligned}
& + \frac{11\langle\bar{s}g_s\sigma Gs\rangle^2}{3072\pi^4} \int dy \left[1 + \frac{s}{2}\delta(s - \tilde{m}_c^2)\right] \\
& - \frac{13\langle\bar{s}g_s\sigma Gs\rangle^2}{27648\pi^4} \int dydz s \delta(s - \overline{m}_c^2) \\
& + \frac{m_s m_c}{3072\pi^8} \int dydz (1-y-z)^3 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s m_c}{12288\pi^8} \int dydz (1-y-z)^4 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s \langle\bar{s}s\rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2)^2 (5s - 2\overline{m}_c^2) \\
& - \frac{m_s \langle\bar{s}s\rangle}{512\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \overline{m}_c^2)^2 (2s - \overline{m}_c^2) \\
& + \frac{11m_s \langle\bar{s}g_s\sigma Gs\rangle}{2048\pi^6} \int dydz (y+z)(1-y-z) (s - \overline{m}_c^2) (2s - \overline{m}_c^2) \\
& + \frac{m_s \langle\bar{s}g_s\sigma Gs\rangle}{1024\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2) (5s - 3\overline{m}_c^2) \\
& - \frac{m_s \langle\bar{s}g_s\sigma Gs\rangle}{2048\pi^6} \int dydz (1-y-z)^2 s (s - \overline{m}_c^2) \\
& + \frac{m_s m_c \langle\bar{s}s\rangle^2}{6\pi^4} \int dydz (s - \overline{m}_c^2) \\
& + \frac{m_s m_c \langle\bar{s}s\rangle^2}{12\pi^4} \int dydz (1-y-z) (s - \overline{m}_c^2) \\
& - \frac{89m_s m_c \langle\bar{s}s\rangle \langle\bar{s}g_s\sigma Gs\rangle}{2304\pi^4} \int dydz \\
& - \frac{199m_s m_c \langle\bar{s}s\rangle \langle\bar{s}g_s\sigma Gs\rangle}{2304\pi^4} \int dy \\
& + \frac{m_s \langle\bar{s}s\rangle^3}{36\pi^2} \int dy \left[1 + \frac{s}{2}\delta(s - \tilde{m}_c^2)\right] \\
& + \frac{m_s m_c \langle\bar{s}g_s\sigma Gs\rangle^2}{64\pi^4} \int dy \left(1 + \frac{7s}{9T^2}\right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle\bar{s}g_s\sigma Gs\rangle^2}{1536\pi^4 T^2} \int dy s \delta(s - \tilde{m}_c^2) , \tag{46}
\end{aligned}$$

$$\begin{aligned}
\rho_{uss}^{11\frac{1}{2},1}(s) &= \frac{1}{61440\pi^8} \int dydz yz(1-y-z)^4 (s - \overline{m}_c^2)^4 (8s - 3\overline{m}_c^2) \\
& - \frac{1}{307200\pi^8} \int dydz yz(1-y-z)^5 (s - \overline{m}_c^2)^4 (9s - 4\overline{m}_c^2) \\
& - \frac{m_c [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{2304\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2)^3 \\
& + \frac{m_c [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{6912\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \overline{m}_c^2)^3 \\
& + \frac{11m_c [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z) (s - \overline{m}_c^2)^2 \\
& - \frac{11m_c [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{24576\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2)^2
\end{aligned}$$

$$\begin{aligned}
& + \frac{\langle \bar{s}s \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{72\pi^4} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{1152\pi^4} \int dydz yz (4s - 3\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{36\pi^2} \int dy \\
& + \frac{11 \langle \bar{s}g_s \sigma Gs \rangle [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{1536\pi^4} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c}{18432\pi^8} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{73728\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s [2\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{192\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{576\pi^6} \int dydz yz(1-y-z)^3 (s - \bar{m}_c^2)^2 (7s - 4\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{192\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2) (3s - 2\bar{m}_c^2) \\
& + \frac{m_s [19\langle \bar{q}g_s \sigma Gq \rangle + 3\langle \bar{s}g_s \sigma Gs \rangle]}{3072\pi^6} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [5\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z) (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [89\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle + 32\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{27648\pi^4} \int dydz (y+z) \\
& - \frac{m_s m_c [231\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 89\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + 256\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c [\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{2304\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{9\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{4608\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{17m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{3456\pi^4} \int dy \left(1 + \frac{8s}{17T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle [3\langle \bar{q}g_s \sigma Gq \rangle + 19\langle \bar{s}g_s \sigma Gs \rangle]}{18432\pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) , \tag{47}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uss}^{11\frac{1}{2},0}(s) &= \frac{1}{245760\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 (7s - 2\bar{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{576\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^3
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{1728\pi^6} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{3072\pi^6} \int dydz (1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{6144\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{4608\pi^4} \int dydz (y+z) (3s - 2\bar{m}_c^2) \\
& + \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle}{2304\pi^4} \int dydz (1-y-z) s \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{9\pi^2} \int dy \\
& + \frac{11\langle \bar{s}g_s\sigma Gs \rangle [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{9216\pi^4} \int dy \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right] \\
& - \frac{13\langle \bar{s}g_s\sigma Gs \rangle [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{82944\pi^4} \int dydz s \delta(s - \bar{m}_c^2) \\
& + \frac{m_s m_c}{4608\pi^8} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{18432\pi^8} \int dydz (1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s [2\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{2304\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{768\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s\sigma Gs \rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{m_s [\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{6144\pi^6} \int dydz (1-y-z)^2 s (s - \bar{m}_c^2) \\
& + \frac{m_s [19\langle \bar{q}g_s\sigma Gq \rangle + 3\langle \bar{s}g_s\sigma Gs \rangle]}{6144\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [5\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{36\pi^4} \int dydz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{36\pi^4} \int dydz (1-y-z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle + 32\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{6912\pi^4} \int dydz \\
& - \frac{m_s m_c [231\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle - 89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 256\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{6912\pi^4} \int dy \\
& + \frac{m_s \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{54\pi^2} \int dy \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right] \\
& + \frac{9m_s m_c \langle \bar{q}g_s\sigma Gq \rangle \langle \bar{s}g_s\sigma Gs \rangle}{864\pi^4} \int dy \left(1 + \frac{8s}{9T^2} \right) \delta(s - \bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s\sigma Gs \rangle [3\langle \bar{q}g_s\sigma Gq \rangle + 19\langle \bar{s}g_s\sigma Gs \rangle]}{13824\pi^4 T^2} \int dy s \delta(s - \bar{m}_c^2) , \tag{48}
\end{aligned}$$

$$\begin{aligned}
\rho_{uus}^{11\frac{1}{2},1}(s) = & \frac{1}{61440\pi^8} \int dydz yz(1-y-z)^4 (s-\overline{m}_c^2)^4 (8s-3\overline{m}_c^2) \\
& - \frac{1}{307200\pi^8} \int dydz yz(1-y-z)^5 (s-\overline{m}_c^2)^4 (9s-4\overline{m}_c^2) \\
& - \frac{m_c [2\langle\bar{q}q\rangle + \langle\bar{s}s\rangle]}{2304\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\overline{m}_c^2)^3 \\
& + \frac{m_c [2\langle\bar{q}q\rangle + \langle\bar{s}s\rangle]}{6912\pi^6} \int dydz (y+z)(1-y-z)^3 (s-\overline{m}_c^2)^3 \\
& + \frac{11m_c [2\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z) (s-\overline{m}_c^2)^2 \\
& - \frac{11m_c [2\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{24576\pi^6} \int dydz (y+z)(1-y-z)^2 (s-\overline{m}_c^2)^2 \\
& + \frac{\langle\bar{q}q\rangle [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{72\pi^4} \int dydz yz(1-y-z) (s-\overline{m}_c^2) (5s-3\overline{m}_c^2) \\
& - \frac{19 [\langle\bar{q}q\rangle\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle + \langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle]}{1152\pi^4} \int dydz yz (4s-3\overline{m}_c^2) \\
& - \frac{m_c\langle\bar{q}q\rangle^2\langle\bar{s}s\rangle}{36\pi^2} \int dy \\
& + \frac{11\langle\bar{q}g_s\sigma Gq\rangle [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{1536\pi^4} \int dy y(1-y) \left[1 + \frac{s}{3}\delta(s-\tilde{m}_c^2)\right] \\
& + \frac{m_s m_c}{36864\pi^8} \int dydz (y+z)(1-y-z)^3 (s-\overline{m}_c^2)^4 \\
& - \frac{m_s m_c}{147456\pi^8} \int dydz (y+z)(1-y-z)^4 (s-\overline{m}_c^2)^4 \\
& - \frac{m_s [4\langle\bar{q}q\rangle - 3\langle\bar{s}s\rangle]}{384\pi^6} \int dydz yz(1-y-z)^2 (s-\overline{m}_c^2)^2 (2s-\overline{m}_c^2) \\
& - \frac{m_s\langle\bar{s}s\rangle}{1152\pi^6} \int dydz yz(1-y-z)^3 (s-\overline{m}_c^2)^2 (7s-4\overline{m}_c^2) \\
& + \frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{384\pi^6} \int dydz yz(1-y-z)^2 (s-\overline{m}_c^2) (3s-2\overline{m}_c^2) \\
& + \frac{m_s [19\langle\bar{q}g_s\sigma Gq\rangle - 8\langle\bar{s}g_s\sigma Gs\rangle]}{3072\pi^6} \int dydz yz(1-y-z) (s-\overline{m}_c^2) (5s-3\overline{m}_c^2) \\
& + \frac{m_s m_c\langle\bar{q}q\rangle [3\langle\bar{q}q\rangle - \langle\bar{s}s\rangle]}{144\pi^4} \int dydz (y+z) (s-\overline{m}_c^2) \\
& + \frac{m_s m_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{144\pi^4} \int dydz (y+z)(1-y-z) (s-\overline{m}_c^2) \\
& - \frac{m_s m_c [32\langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle + 57\langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle]}{27648\pi^4} \int dydz (y+z) \\
& - \frac{m_s m_c [288\langle\bar{q}q\rangle\langle\bar{q}g_s\sigma Gq\rangle - 57\langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle - 32\langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c\langle\bar{q}q\rangle\langle\bar{q}g_s\sigma Gq\rangle}{2304\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y}\right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{18\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle^2}{9216\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle [144 \langle \bar{q}g_s \sigma Gq \rangle - 19 \langle \bar{s}g_s \sigma Gs \rangle]}{55296\pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2), \quad (49)
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uus}^{11\frac{1}{2},0}(s) = & \frac{1}{245760\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \overline{m}_c^2)^4 (7s - 2\overline{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \overline{m}_c^2)^4 (8s - 3\overline{m}_c^2) \\
& - \frac{m_c [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{576\pi^6} \int dydz (1-y-z)^2 (s - \overline{m}_c^2)^3 \\
& + \frac{m_c [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{1728\pi^6} \int dydz (1-y-z)^3 (s - \overline{m}_c^2)^3 \\
& + \frac{11m_c [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{3072\pi^6} \int dydz (1-y-z) (s - \overline{m}_c^2)^2 \\
& - \frac{11m_c [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{6144\pi^6} \int dydz (1-y-z)^2 (s - \overline{m}_c^2)^2 \\
& + \frac{\langle \bar{q}q \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (y+z)(1-y-z) (s - \overline{m}_c^2) (2s - \overline{m}_c^2) \\
& - \frac{19 [\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{4608\pi^4} \int dydz (y+z) (3s - 2\overline{m}_c^2) \\
& + \frac{\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{2304\pi^4} \int dydz (1-y-z) s \\
& - \frac{m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{9\pi^2} \int dy \\
& + \frac{11 \langle \bar{q}g_s \sigma Gq \rangle [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{9216\pi^4} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{13 \langle \bar{q}g_s \sigma Gq \rangle [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{82944\pi^4} \int dydz s \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c}{9216\pi^8} \int dydz (1-y-z)^3 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s m_c}{36864\pi^8} \int dydz (1-y-z)^4 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s [4\langle \bar{q}q \rangle - 3\langle \bar{s}s \rangle]}{4608\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2)^2 (5s - 2\overline{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \overline{m}_c^2)^2 (2s - \overline{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{3072\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2) (5s - 3\overline{m}_c^2) \\
& - \frac{m_s \langle \bar{q}g_s \sigma Gq \rangle}{6144\pi^6} \int dydz (1-y-z)^2 s (s - \overline{m}_c^2) \\
& + \frac{m_s [19\langle \bar{q}g_s \sigma Gq \rangle - 8\langle \bar{s}g_s \sigma Gs \rangle]}{6144\pi^6} \int dydz (y+z)(1-y-z) (s - \overline{m}_c^2) (2s - \overline{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s m_c \langle \bar{q}q \rangle [3\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{36\pi^4} \int dydz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{36\pi^4} \int dydz (1 - y - z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [32\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{6912\pi^4} \int dydz \\
& - \frac{m_s m_c [288\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle - 57\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 32\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle]}{6912\pi^4} \int dy \\
& + \frac{m_s \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{108\pi^2} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle^2}{192\pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{19m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{13824\pi^4 T^2} \int dy s \delta(s - \tilde{m}_c^2) , \tag{50}
\end{aligned}$$

$$\rho_{uu\bar{u}}^{11\frac{1}{2},1}(s) = \rho_{ss\bar{s}}^{11\frac{1}{2},1}(s) \big|_{m_s \rightarrow 0, \langle \bar{s}s \rangle \rightarrow \langle \bar{q}q \rangle, \langle \bar{s}g_s \sigma Gs \rangle \rightarrow \langle \bar{q}g_s \sigma Gq \rangle} , \tag{51}$$

$$\tilde{\rho}_{uu\bar{u}}^{11\frac{1}{2},0}(s) = \tilde{\rho}_{ss\bar{s}}^{11\frac{1}{2},0}(s) \big|_{m_s \rightarrow 0, \langle \bar{s}s \rangle \rightarrow \langle \bar{q}q \rangle, \langle \bar{s}g_s \sigma Gs \rangle \rightarrow \langle \bar{q}g_s \sigma Gq \rangle} , \tag{52}$$

$$\begin{aligned}
\rho_{ss\bar{s}}^{10\frac{1}{2},1}(s) = & \frac{1}{245760\pi^8} \int dydz yz(1-y-z)^4 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dydz yz(1-y-z)^5 (s - \bar{m}_c^2)^4 (9s - 4\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s}s \rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c \langle \bar{s}s \rangle}{4608\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{8192\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{16384\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& - \frac{7m_c \langle \bar{s}g_s \sigma Gs \rangle}{16384\pi^6} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{m_c \langle \bar{s}g_s \sigma Gs \rangle}{8192\pi^6} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle^2}{96\pi^4} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{19\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{1536\pi^4} \int dydz yz (4s - 3\bar{m}_c^2) \\
& - \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{3072\pi^4} \int dydz (y+z)(1-y-z) (2s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{s}s \rangle^3}{72\pi^2} \int dy \\
& + \frac{13\langle \bar{s}g_s \sigma Gs \rangle^2}{110592\pi^4} \int dydz (y+z) [1 + s \delta(s - \bar{m}_c^2)]
\end{aligned}$$

$$\begin{aligned}
& + \frac{11\langle\bar{s}g_s\sigma Gs\rangle^2}{2048\pi^4} \int dy y(1-y) \left[1 + \frac{s}{3}\delta(s - \tilde{m}_c^2)\right] \\
& + \frac{m_s m_c}{24576\pi^8} \int dydz (y+z)(1-y-z)^3 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s m_c}{98304\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \overline{m}_c^2)^4 \\
& - \frac{m_s\langle\bar{s}s\rangle}{512\pi^6} \int dydz yz(1-y-z)^2 (s - \overline{m}_c^2)^2 (2s - \overline{m}_c^2) \\
& - \frac{m_s\langle\bar{s}s\rangle}{1536\pi^6} \int dydz yz(1-y-z)^3 (s - \overline{m}_c^2)^2 (7s - 4\overline{m}_c^2) \\
& + \frac{11m_s\langle\bar{s}g_s\sigma Gs\rangle}{4096\pi^6} \int dydz yz(1-y-z) (s - \overline{m}_c^2) (5s - 3\overline{m}_c^2) \\
& + \frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{512\pi^6} \int dydz yz(1-y-z)^2 (s - \overline{m}_c^2) (3s - 2\overline{m}_c^2) \\
& + \frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{16384\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \overline{m}_c^2) (3s - \overline{m}_c^2) \\
& + \frac{m_s m_c\langle\bar{s}s\rangle^2}{48\pi^4} \int dydz (y+z) (s - \overline{m}_c^2) \\
& + \frac{m_s m_c\langle\bar{s}s\rangle^2}{96\pi^4} \int dydz (y+z)(1-y-z) (s - \overline{m}_c^2) \\
& - \frac{89m_s m_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{18432\pi^4} \int dydz (y+z) \\
& - \frac{199m_s m_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{18432\pi^4} \int dy \\
& + \frac{3m_s m_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{1024\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y}\right) \\
& + \frac{m_s m_c\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{512\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y}\right) (1-y-z) \\
& + \frac{m_s\langle\bar{s}s\rangle^3}{24\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3}\delta(s - \tilde{m}_c^2)\right] \\
& + \frac{m_s m_c\langle\bar{s}g_s\sigma Gs\rangle^2}{4608\pi^4} \int dy \left(16 + \frac{7s}{T^2}\right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c\langle\bar{s}g_s\sigma Gs\rangle^2}{3072\pi^4} \int dydz \left(\frac{z}{y} + \frac{y}{z}\right) \delta(s - \overline{m}_c^2) \\
& - \frac{17m_s m_c\langle\bar{s}g_s\sigma Gs\rangle^2}{18432\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y}\right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c\langle\bar{s}g_s\sigma Gs\rangle^2}{12288\pi^4} \int dy \left(1 + \frac{s}{T^2}\right) \delta(s - \tilde{m}_c^2) , \tag{53}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{ss\bar{s}\bar{s}}^{10\frac{1}{2},0}(s) &= \frac{1}{491520\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \overline{m}_c^2)^4 (7s - 2\overline{m}_c^2) \\
& - \frac{1}{2457600\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \overline{m}_c^2)^4 (8s - 3\overline{m}_c^2) \\
& - \frac{m_c\langle\bar{s}s\rangle}{768\pi^6} \int dydz (1-y-z)^2 (s - \overline{m}_c^2)^3
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_c \langle \bar{s}s \rangle}{2304\pi^6} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{4096\pi^6} \int dydz (1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c \langle \bar{s}g_s \sigma Gs \rangle}{8192\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& - \frac{7m_c \langle \bar{s}g_s \sigma Gs \rangle}{16384\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{m_c \langle \bar{s}g_s \sigma Gs \rangle}{8192\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle^2}{96\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{3072\pi^4} \int dydz (y+z) (3s - 2\bar{m}_c^2) \\
& - \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{1536\pi^4} \int dydz (1-y-z)s \\
& - \frac{m_c \langle \bar{s}s \rangle^3}{36\pi^2} \int dy \\
& + \frac{11\langle \bar{s}g_s \sigma Gs \rangle^2}{6144\pi^4} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{13\langle \bar{s}g_s \sigma Gs \rangle^2}{55296\pi^4} \int dydz s \delta(s - \bar{m}_c^2) \\
& + \frac{m_s m_c}{12288\pi^8} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{49152\pi^8} \int dydz (1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s \langle \bar{s}s \rangle}{3072\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{1024\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& + \frac{11m_s \langle \bar{s}g_s \sigma Gs \rangle}{4096\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{2048\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{4096\pi^6} \int dydz (1-y-z)^2 s (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle^2}{24\pi^4} \int dydz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle^2}{48\pi^4} \int dydz (1-y-z) (s - \bar{m}_c^2) \\
& - \frac{89m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{9216\pi^4} \int dydz \\
& - \frac{199m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{9216\pi^4} \int dy \\
& + \frac{3m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{1024\pi^4} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s m_c \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{512\pi^4} \int dy dz \left(\frac{1}{y} + \frac{1}{z} \right) (1 - y - z) \\
& + \frac{m_s \langle \bar{s}s \rangle^3}{72\pi^2} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{256\pi^4} \int dy \left(1 + \frac{7s}{9T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{3072\pi^4} \int dy dz \left(\frac{1}{y} + \frac{1}{z} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{17m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{18432\pi^4} \int dy \left(\frac{1}{y} + \frac{1}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s}g_s \sigma Gs \rangle^2}{6144\pi^4 T^2} \int dy s \delta(s - \tilde{m}_c^2) , \tag{54}
\end{aligned}$$

$$\begin{aligned}
\rho_{us\bar{s}}^{10\frac{1}{2},1}(s) = & \frac{1}{245760\pi^8} \int dy dz yz(1-y-z)^4 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{1}{1228800\pi^8} \int dy dz yz(1-y-z)^5 (s - \bar{m}_c^2)^4 (9s - 4\bar{m}_c^2) \\
& - \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{4608\pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{13824\pi^6} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{24576\pi^6} \int dy dz (y+z)(1-y-z) (s - \bar{m}_c^2)^2 \\
& + \frac{11m_c [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{49152\pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& - \frac{7m_c [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{49152\pi^6} \int dy dz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{m_c [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{24576\pi^6} \int dy dz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{288\pi^4} \int dy dz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{4608\pi^4} \int dy dz yz (4s - 3\bar{m}_c^2) \\
& - \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{9216\pi^4} \int dy dz (y+z)(1-y-z) (2s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{72\pi^2} \int dy \\
& + \frac{11 \langle \bar{s}g_s \sigma Gs \rangle [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{6144\pi^4} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{13 \langle \bar{s}g_s \sigma Gs \rangle [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{331776\pi^4} \int dy dz (y+z) \left[1 + s \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c}{36864\pi^8} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{147456\pi^8} \int dy dz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4
\end{aligned}$$

$$\begin{aligned}
& -\frac{m_s [2\langle\bar{q}q\rangle - \langle\bar{s}s\rangle]}{768\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& -\frac{m_s \langle\bar{s}s\rangle}{2304\pi^6} \int dydz yz(1-y-z)^3 (s - \bar{m}_c^2)^2 (7s - 4\bar{m}_c^2) \\
& +\frac{m_s \langle\bar{s}g_s\sigma Gs\rangle}{768\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2) (3s - 2\bar{m}_c^2) \\
& +\frac{m_s [19\langle\bar{q}g_s\sigma Gq\rangle + 3\langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& +\frac{m_s [\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{49152\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (3s - \bar{m}_c^2) \\
& +\frac{m_s m_c \langle\bar{s}s\rangle [5\langle\bar{q}q\rangle - \langle\bar{s}s\rangle]}{288\pi^4} \int dydz (y+z) (s - \bar{m}_c^2) \\
& +\frac{m_s m_c \langle\bar{s}s\rangle [\langle\bar{q}q\rangle + \langle\bar{s}s\rangle]}{288\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) \\
& -\frac{m_s m_c [89\langle\bar{s}s\rangle \langle\bar{s}g_s\sigma Gs\rangle + 32\langle\bar{q}q\rangle \langle\bar{s}g_s\sigma Gs\rangle + 57\langle\bar{s}s\rangle \langle\bar{q}g_s\sigma Gq\rangle]}{55296\pi^4} \int dydz (y+z) \\
& -\frac{m_s m_c [231\langle\bar{s}s\rangle \langle\bar{q}g_s\sigma Gq\rangle + 256\langle\bar{q}q\rangle \langle\bar{s}g_s\sigma Gs\rangle - 89\langle\bar{s}s\rangle \langle\bar{s}g_s\sigma Gs\rangle]}{55296\pi^4} \int dy \\
& +\frac{m_s m_c \langle\bar{s}s\rangle [\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{1536\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y}\right) (1-y-z) \\
& +\frac{m_s m_c [9\langle\bar{s}s\rangle \langle\bar{q}g_s\sigma Gq\rangle - 7\langle\bar{s}s\rangle \langle\bar{s}g_s\sigma Gs\rangle + 16\langle\bar{q}q\rangle \langle\bar{s}g_s\sigma Gs\rangle]}{9216\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y}\right) \\
& +\frac{m_s \langle\bar{q}q\rangle \langle\bar{s}s\rangle^2}{36\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2)\right] \\
& -\frac{m_s m_c \langle\bar{s}g_s\sigma Gs\rangle [\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{9216\pi^4} \int dydz \left(\frac{z}{y} + \frac{y}{z}\right) \delta(s - \bar{m}_c^2) \\
& -\frac{m_s m_c \langle\bar{s}g_s\sigma Gs\rangle [41\langle\bar{q}g_s\sigma Gq\rangle - 7\langle\bar{s}g_s\sigma Gs\rangle]}{55296\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y}\right) \delta(s - \tilde{m}_c^2) \\
& -\frac{m_s m_c \langle\bar{s}g_s\sigma Gs\rangle [19\langle\bar{s}g_s\sigma Gs\rangle - 125\langle\bar{q}g_s\sigma Gq\rangle]}{55296\pi^4} \int dy \left(1 + \frac{s}{2T^2}\right) \delta(s - \tilde{m}_c^2) \\
& +\frac{19m_s m_c \langle\bar{s}g_s\sigma Gs\rangle [\langle\bar{q}g_s\sigma Gq\rangle + \langle\bar{s}g_s\sigma Gs\rangle]}{110592\pi^4} \int dy \delta(s - \tilde{m}_c^2), \tag{55}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uss}^{10\frac{1}{2},0}(s) &= \frac{1}{491520\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 (7s - 2\bar{m}_c^2) \\
& -\frac{1}{2457600\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& -\frac{m_c [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{2304\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& +\frac{m_c [\langle\bar{q}q\rangle + 2\langle\bar{s}s\rangle]}{6912\pi^6} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& +\frac{11m_c [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{12288\pi^6} \int dydz (1-y-z) (s - \bar{m}_c^2)^2 \\
& -\frac{11m_c [\langle\bar{q}g_s\sigma Gq\rangle + 2\langle\bar{s}g_s\sigma Gs\rangle]}{24576\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^2
\end{aligned}$$

$$\begin{aligned}
& - \frac{7m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{49152\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{m_c [\langle \bar{q}g_s\sigma Gq \rangle + 2\langle \bar{s}g_s\sigma Gs \rangle]}{24576\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{s}s \rangle [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{288\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19 [\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{9216\pi^4} \int dydz (y+z) (3s - 2\bar{m}_c^2) \\
& - \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle}{4608\pi^4} \int dydz (1-y-z)s \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{36\pi^2} \int dy \\
& + \frac{13 \langle \bar{s}g_s\sigma Gs \rangle [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{165888\pi^4} \int dydz s \delta(s - \bar{m}_c^2) \\
& + \frac{11 \langle \bar{s}g_s\sigma Gs \rangle [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{18432\pi^4} \int dy \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right] \\
& + \frac{m_s m_c}{18432\pi^8} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{73728\pi^8} \int dydz (1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s [2\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{4608\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{1536\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s\sigma Gs \rangle}{3072\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s [\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{12288\pi^6} \int dydz (1-y-z)^2 s (s - \bar{m}_c^2) \\
& + \frac{m_s [19\langle \bar{q}g_s\sigma Gq \rangle + 3\langle \bar{s}g_s\sigma Gs \rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [5\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (1-y-z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 32\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{27648\pi^4} \int dydz \\
& - \frac{m_s m_c [231\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle + 256\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle - 89\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c \langle \bar{s}s \rangle [\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{1536\pi^4} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z) \\
& + \frac{m_s m_c [9\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle - 7\langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle + 16\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle]}{9216\pi^4} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{108\pi^2} \int dy \left[1 + \frac{s}{2} \delta(s - \bar{m}_c^2) \right]
\end{aligned}$$

$$\begin{aligned}
& - \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle [\langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{9216 \pi^4} \int dy dz \left(\frac{1}{y} + \frac{1}{z} \right) \delta(s - \bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle [41 \langle \bar{q} g_s \sigma G q \rangle - 7 \langle \bar{s} g_s \sigma G s \rangle]}{55296 \pi^4} \int dy \left(\frac{1}{y} + \frac{1}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle [19 \langle \bar{s} g_s \sigma G s \rangle - 125 \langle \bar{q} g_s \sigma G q \rangle]}{55296 \pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{19 m_s m_c \langle \bar{s} g_s \sigma G s \rangle [\langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{55296 \pi^4} \int dy \delta(s - \tilde{m}_c^2) , \tag{56}
\end{aligned}$$

$$\begin{aligned}
\rho_{uus}^{10\frac{1}{2},1}(s) = & \frac{1}{245760 \pi^8} \int dy dz y z (1-y-z)^4 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{1}{1228800 \pi^8} \int dy dz y z (1-y-z)^5 (s - \bar{m}_c^2)^4 (9s - 4\bar{m}_c^2) \\
& - \frac{m_c [2 \langle \bar{q} q \rangle + \langle \bar{s} s \rangle]}{4608 \pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c [2 \langle \bar{q} q \rangle + \langle \bar{s} s \rangle]}{13824 \pi^6} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11 m_c [2 \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{24576 \pi^6} \int dy dz (y+z)(1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11 m_c [2 \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{49152 \pi^6} \int dy dz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& - \frac{7 m_c [2 \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{49152 \pi^6} \int dy dz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& + \frac{m_c [2 \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{s} g_s \sigma G s \rangle]}{24576 \pi^6} \int dy dz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{q} q \rangle [\langle \bar{q} q \rangle + 2 \langle \bar{s} s \rangle]}{288 \pi^4} \int dy dz y z (1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& - \frac{\langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + \langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle}{9216 \pi^4} \int dy dz (y+z)(1-y-z) (2s - \bar{m}_c^2) \\
& - \frac{19 [\langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle + \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + \langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle]}{4608 \pi^4} \int dy dz y z (4s - 3\bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q} q \rangle^2 \langle \bar{s} s \rangle}{72 \pi^2} \int dy \\
& + \frac{11 \langle \bar{q} g_s \sigma G q \rangle [\langle \bar{q} g_s \sigma G q \rangle + 2 \langle \bar{s} g_s \sigma G s \rangle]}{6144 \pi^4} \int dy y (1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{13 \langle \bar{q} g_s \sigma G q \rangle [\langle \bar{q} g_s \sigma G q \rangle + 2 \langle \bar{s} g_s \sigma G s \rangle]}{331776 \pi^4} \int dy dz (y+z) \left[1 + s \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c}{73728 \pi^8} \int dy dz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{294912 \pi^8} \int dy dz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s [4 \langle \bar{q} q \rangle - 3 \langle \bar{s} s \rangle]}{1536 \pi^6} \int dy dz y z (1-y-z)^2 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s} s \rangle}{4608 \pi^6} \int dy dz y z (1-y-z)^3 (s - \bar{m}_c^2)^2 (7s - 4\bar{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s [19\langle \bar{q}g_s\sigma Gq \rangle - 8\langle \bar{s}g_s\sigma Gs \rangle]}{12288\pi^6} \int dydz yz(1-y-z) (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s\sigma Gs \rangle}{1536\pi^6} \int dydz yz(1-y-z)^2 (s - \bar{m}_c^2) (3s - 2\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{q}g_s\sigma Gq \rangle}{49152\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (3s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle [3\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{288\pi^4} \int dydz (y+z) (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{288\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [32\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{55296\pi^4} \int dydz (y+z) \\
& - \frac{m_s m_c [288\langle \bar{q}q \rangle \langle \bar{q}g_s\sigma Gq \rangle - 32\langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle - 57\langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle]}{55296\pi^4} \int dy \\
& + \frac{m_s m_c \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle}{1536\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) (1-y-z) \\
& + \frac{m_s m_c [16\langle \bar{q}q \rangle - 7\langle \bar{s}s \rangle] \langle \bar{q}g_s\sigma Gq \rangle}{9216\pi^4} \int dydz \left(\frac{y}{z} + \frac{z}{y} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{72\pi^2} \int dy y(1-y) \left[1 + \frac{s}{3} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle \langle \bar{s}g_s\sigma Gs \rangle}{6912\pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle \langle \bar{s}g_s\sigma Gs \rangle}{9216\pi^4} \int dydz \left(\frac{z}{y} + \frac{y}{z} \right) \delta(s - \bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle [24\langle \bar{q}g_s\sigma Gq \rangle - 7\langle \bar{s}g_s\sigma Gs \rangle]}{55296\pi^4} \int dy \left(\frac{1-y}{y} + \frac{y}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle [24\langle \bar{q}g_s\sigma Gq \rangle - \langle \bar{s}g_s\sigma Gs \rangle]}{18432\pi^4} \int dy \left(1 + \frac{s}{2T^2} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle \langle \bar{s}g_s\sigma Gs \rangle}{36864\pi^4} \int dy \delta(s - \tilde{m}_c^2) , \tag{57}
\end{aligned}$$

$$\begin{aligned}
\tilde{\rho}_{uus}^{10\frac{1}{2},0}(s) &= \frac{1}{491520\pi^8} \int dydz (y+z)(1-y-z)^4 (s - \bar{m}_c^2)^4 (7s - 2\bar{m}_c^2) \\
& - \frac{1}{2457600\pi^8} \int dydz (y+z)(1-y-z)^5 (s - \bar{m}_c^2)^4 (8s - 3\bar{m}_c^2) \\
& - \frac{m_c [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{2304\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^3 \\
& + \frac{m_c [2\langle \bar{q}q \rangle + \langle \bar{s}s \rangle]}{6912\pi^6} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^3 \\
& + \frac{11m_c [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{12288\pi^6} \int dydz (1-y-z) (s - \bar{m}_c^2)^2 \\
& - \frac{11m_c [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{24576\pi^6} \int dydz (1-y-z)^2 (s - \bar{m}_c^2)^2 \\
& - \frac{7m_c [2\langle \bar{q}g_s\sigma Gq \rangle + \langle \bar{s}g_s\sigma Gs \rangle]}{49152\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^2 (s - \bar{m}_c^2)^2
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_c [2\langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{s}g_s \sigma Gs \rangle]}{24576\pi^6} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z)^3 (s - \bar{m}_c^2)^2 \\
& + \frac{\langle \bar{q}q \rangle [\langle \bar{q}q \rangle + 2\langle \bar{s}s \rangle]}{288\pi^4} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& - \frac{19 [\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{9216\pi^4} \int dydz (y+z) (3s - 2\bar{m}_c^2) \\
& - \frac{\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle + \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{4608\pi^4} \int dydz (1-y-z)s \\
& - \frac{m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{36\pi^2} \int dy \\
& + \frac{11 \langle \bar{q}g_s \sigma Gq \rangle [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{18432\pi^4} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{13 \langle \bar{q}g_s \sigma Gq \rangle [\langle \bar{q}g_s \sigma Gq \rangle + 2\langle \bar{s}g_s \sigma Gs \rangle]}{165888\pi^4} \int dydz s \delta(s - \bar{m}_c^2) \\
& + \frac{m_s m_c}{36864\pi^8} \int dydz (1-y-z)^3 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s m_c}{147456\pi^8} \int dydz (1-y-z)^4 (s - \bar{m}_c^2)^4 \\
& - \frac{m_s [4\langle \bar{q}q \rangle - 3\langle \bar{s}s \rangle]}{9216\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2)^2 (5s - 2\bar{m}_c^2) \\
& - \frac{m_s \langle \bar{s}s \rangle}{3072\pi^6} \int dydz (y+z)(1-y-z)^3 (s - \bar{m}_c^2)^2 (2s - \bar{m}_c^2) \\
& + \frac{m_s [19\langle \bar{q}g_s \sigma Gq \rangle - 8\langle \bar{s}g_s \sigma Gs \rangle]}{12288\pi^6} \int dydz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{6144\pi^6} \int dydz (y+z)(1-y-z)^2 (s - \bar{m}_c^2) (5s - 3\bar{m}_c^2) \\
& + \frac{m_s \langle \bar{q}g_s \sigma Gq \rangle}{12288\pi^6} \int dydz (1-y-z)^2 s (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle [3\langle \bar{q}q \rangle - \langle \bar{s}s \rangle]}{144\pi^4} \int dydz (s - \bar{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{144\pi^4} \int dydz (1-y-z) (s - \bar{m}_c^2) \\
& - \frac{m_s m_c [32\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 57\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{27648\pi^4} \int dydz \\
& - \frac{m_s m_c [288\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle - 32\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle - 57\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle]}{27648\pi^4} \int dy \\
& + \frac{m_s m_c \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{1536\pi^4} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) (1-y-z) \\
& + \frac{m_s m_c [16\langle \bar{q}q \rangle - 7\langle \bar{s}s \rangle] \langle \bar{q}g_s \sigma Gq \rangle}{9216\pi^4} \int dydz \left(\frac{1}{y} + \frac{1}{z} \right) \\
& + \frac{m_s \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{216\pi^2} \int dy \left[1 + \frac{s}{2} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{3456\pi^4 T^2} \int dy s \delta(s - \tilde{m}_c^2)
\end{aligned}$$

$$\begin{aligned}
& - \frac{m_s m_c \langle \bar{q} g_s \sigma G q \rangle \langle \bar{s} g_s \sigma G s \rangle}{9216 \pi^4} \int dy dz \left(\frac{1}{y} + \frac{1}{z} \right) \delta(s - \bar{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{q} g_s \sigma G q \rangle [24 \langle \bar{q} g_s \sigma G q \rangle - 7 \langle \bar{s} g_s \sigma G s \rangle]}{55296 \pi^4} \int dy \left(\frac{1}{y} + \frac{1}{1-y} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q} g_s \sigma G q \rangle [24 \langle \bar{q} g_s \sigma G q \rangle - \langle \bar{s} g_s \sigma G s \rangle]}{18432 \pi^4} \int dy \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q} g_s \sigma G q \rangle \langle \bar{s} g_s \sigma G s \rangle}{18432 \pi^4} \int dy \delta(s - \tilde{m}_c^2), \tag{58}
\end{aligned}$$

$$\rho_{uuu}^{10\frac{1}{2},1}(s) = \rho_{sss}^{10\frac{1}{2},1}(s) \big|_{m_s \rightarrow 0, \langle \bar{s}s \rangle \rightarrow \langle \bar{q}q \rangle, \langle \bar{s}g_s \sigma G s \rangle \rightarrow \langle \bar{q}g_s \sigma G q \rangle}, \tag{59}$$

$$\tilde{\rho}_{uuu}^{10\frac{1}{2},0}(s) = \tilde{\rho}_{sss}^{10\frac{1}{2},0}(s) \big|_{m_s \rightarrow 0, \langle \bar{s}s \rangle \rightarrow \langle \bar{q}q \rangle, \langle \bar{s}g_s \sigma G s \rangle \rightarrow \langle \bar{q}g_s \sigma G q \rangle}, \tag{60}$$

where $\int dy dz = \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz$, $\int dy = \int_{y_i}^{y_f} dy$, $y_f = \frac{1+\sqrt{1-4m_c^2/s}}{2}$, $y_i = \frac{1-\sqrt{1-4m_c^2/s}}{2}$, $z_i = \frac{ym_c^2}{ys-\tilde{m}_c^2}$, $\bar{m}_c^2 = \frac{(y+z)m_c^2}{yz}$, $\tilde{m}_c^2 = \frac{m_c^2}{y(1-y)}$, $\int_{y_i}^{y_f} dy \rightarrow \int_0^1 dy$, $\int_{z_i}^{1-y} dz \rightarrow \int_0^{1-y} dz$, when the $\delta(s - \bar{m}_c^2)$ and $\delta(s - \tilde{m}_c^2)$ appear.

Acknowledgements

This work is supported by National Natural Science Foundation, Grant Number 11375063, and Natural Science Foundation of Hebei province, Grant Number A2014502017.

References

- [1] R. Aaij et al, Phys. Rev. Lett. **115** (2015) 072001.
- [2] R. Aaij et al, arXiv:1604.05708.
- [3] R. Aaij et al, arXiv:1606.06999.
- [4] R. Chen, X. Liu, X. Q. Li and S. L. Zhu, Phys. Rev. Lett. **115** (2015) 132002; H. X. Chen, W. Chen, X. Liu, T. G. Steele and S. L. Zhu, Phys. Rev. Lett. **115** (2015) 172001; U. G. Meissner and J. A. Oller, Phys. Lett. **B751** (2015) 59; C. W. Xiao and U. G. Meissner, Phys. Rev. **D92** (2015) 114002; N. N. Scoccola, D. O. Riska and M. Rho, Phys. Rev. **D92** (2015) 051501; L. Roca, J. Nieves and E. Oset, Phys. Rev. **D92** (2015) 094003; T. J. Burns, Eur. Phys. J. **A51** (2015) 152; H. X. Chen, L. S. Geng, W. H. Liang, E. Oset, E. Wang and J. J. Xie, arXiv:1510.01803; G. Yang and J. Ping, arXiv:1511.09053; J. He, Phys. Lett. **B753** (2016) 547; E. Wang, H. X. Chen, L. S. Geng, D. M. Li and E. Oset, Phys. Rev. **D93** (2016) 094001; G. J. Wang, L. Ma, X. Liu and S. L. Zhu, Phys. Rev. **D93** (2016) 034031.
- [5] A. Mironov and A. Morozov, JETP Lett. **102** (2015) 271.
- [6] R. F. Lebed, Phys. Rev. **D92** (2015) 114030; R. F. Lebed, Phys. Lett. **B749** (2015) 454; R. Zhu and C. F. Qiao, Phys. Lett. **B756** (2016) 259.
- [7] L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. **B749** (2015) 289.
- [8] V. V. Anisovich, M. A. Matveev, J. Nyiri, A. V. Sarantsev and A. N. Semenova, arXiv:1507.07652; R. Ghosh, A. Bhattacharya and B. Chakrabarti, arXiv:1508.00356; V. V. Anisovich, M. A. Matveev, J. Nyiri, A. V. Sarantsev and A. N. Semenova, Int. J. Mod. Phys. **A30** (2015) 1550190; L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. **B750** (2015) 37; H.

- Y. Cheng and C. K. Chua, Phys. Rev. **D92** (2015) 096009; V. V. Anisovich, M. A. Matveev, A. V. Sarantsev and A. N. Semenova, Mod. Phys. Lett. **A30** (2015) 1550212; G. N. Li, M. He and X. G. He, JHEP **1512** (2015) 128.
- [9] Z. G. Wang, Eur. Phys. J. **C76** (2016) 70.
- [10] Z. G. Wang and T. Huang, Eur. Phys. J. **C76** (2016) 43.
- [11] Z. G. Wang, Eur. Phys. J. **C76** (2016) 142.
- [12] F. K. Guo, U. G. Meissner, W. Wang and Z. Yang, Phys. Rev. **D92** (2015) 071502; M. Mikhasenko, arXiv:1507.06552; X. H. Liu, Q. Wang and Q. Zhao, Phys. Lett. **B757** (2016) 231.
- [13] Q. Wang, X. H. Liu and Q. Zhao, Phys. Rev. **D92** (2015) 034022; V. Kubarovsky and M. B. Voloshin, Phys. Rev. **D92** (2015) 031502; Q. F. Lu, X. Y. Wang, J. J. Xie, X. R. Chen and Y. B. Dong, Phys. Rev. **D93** (2016) 034009; M. Karliner and J. L. Rosner, Phys. Lett. **B752** (2016) 329.
- [14] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385, 448.
- [15] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.
- [16] P. Colangelo and A. Khodjamirian, hep-ph/0010175.
- [17] S. Narison, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17** (2002) 1.
- [18] A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. **D12** (1975) 147; T. DeGrand, R. L. Jaffe, K. Johnson and J. E. Kiskis, Phys. Rev. **D12** (1975) 2060; R. L. Jaffe, Phys. Rept. **409** (2005) 1.
- [19] Z. G. Wang, Eur. Phys. J. **C71** (2011) 1524; R. T. Kleiv, T. G. Steele and A. Zhang, Phys. Rev. **D87** (2013) 125018.
- [20] Z. G. Wang, Commun. Theor. Phys. **59** (2013) 451.
- [21] Z. G. Wang and T. Huang, Phys. Rev. **D89** (2014) 054019; Z. G. Wang, Eur. Phys. J. **C74** (2014) 2874; Z. G. Wang and T. Huang, Nucl. Phys. **A930** (2014) 63.
- [22] Y. Chung, H. G. Dosch, M. Kremer and D. Schall, Nucl. Phys. **B197** (1982) 55.
- [23] E. Bagan, M. Chabab, H. G. Dosch and S. Narison, Phys. Lett. **B301**, 243 (1993).
- [24] D. Jido, N. Kodama and M. Oka, Phys. Rev. **D54** (1996) 4532.
- [25] Z. G. Wang, Phys. Lett. **B685** (2010) 59; Z. G. Wang, Eur. Phys. J. **A45** (2010) 267; Z. G. Wang, Eur. Phys. J. **C68** (2010) 459; Z. G. Wang, Eur. Phys. J. **A47** (2011) 81; Z. G. Wang, Commun. Theor. Phys. **58** (2012) 723.
- [26] K. A. Olive et al, Chin. Phys. **C38** (2014) 090001.
- [27] Z. G. Wang, Eur. Phys. J. **C68** (2010) 479.
- [28] S. J. Brodsky, D. S. Hwang and R. F. Lebed, Phys. Rev. Lett. **113** (2014) 112001.
- [29] M. Neubert, Phys. Rept. **245** (1994) 259.